


WHO Child Growth Standards

Length/height-for-age, weight-for-age, weight-for-length,
weight-for-height and body mass index-for-age

Methods and development



World Health
Organization

WHO Child Growth Standards

**Length/height-for-age, weight-for-age, weight-for-length,
weight-for-height and body mass index-for-age**

Methods and development



**World Health
Organization**

**Department of Nutrition for
Health and Development**

WHO Library Cataloguing-in-Publication Data

WHO child growth standards : length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age : methods and development.

Coordinating team: Mercedes de Onis ... [et al.].

1. Anthropometry. 2. Anthropometry - methods. 3. Body weights and measures - standards. 4. Child development. 5. Growth. 6. Reference standards. 7. Nutrition assessment. I. de Onis, Mercedes. II. World Health Organization. III. Title: World Health Organization child growth standards.

ISBN 92 4 154693 X

(NLM classification: WS 103)

© World Health Organization 2006

All rights reserved. Publications of the World Health Organization can be obtained from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel: +41 22 791 3264; fax: +41 22 791 4857; email: bookorders@who.int). Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to WHO Press, at the above address (fax: +41 22 791 4806; email: permissions@who.int).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either express or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Printed in France

Members of the WHO Multicentre Growth Reference Study Group

Coordinating Team

Mercedes de Onis [Study Coordinator], Adelheid Onyango, Elaine Borghi, Amani Siyam, Alain Pinol (Department of Nutrition for Health and Development, World Health Organization).

Executive Committee

Cutberto Garza [Chair], Mercedes de Onis, Jose Martines, Reynaldo Martorell, Cesar G. Victora (up to October 2002), Maharaj K. Bhan (from November 2002).

Steering Committee

Coordinating Centre (WHO, Geneva): Mercedes de Onis, Jose Martines, Adelheid Onyango, Alain Pinol.

Investigators (by country): Cesar G. Victora and Cora Luiza Araújo (Brazil), Anna Lartey and William B. Owusu (Ghana), Maharaj K. Bhan and Nita Bhandari (India), Kaare R. Norum and Gunn-Elin Aa. Bjoerneboe (Norway), Ali Jaffer Mohamed (Oman), Kathryn G. Dewey (USA).

United Nations Agency Representatives: Cutberto Garza (UNU), Krishna Belbase (UNICEF).

Advisory Group

Maureen Black, Wm. Cameron Chumlea, Tim Cole, Edward Frongillo, Laurence Grummer-Strawn, Reynaldo Martorell, Roger Shrimpton, Jan Van den Broeck. For the work presented in this document, Huiqi Pan, Robert Rigby, Mikis Stasinopoulos and Stef van Buuren, participated in an advisory capacity.

Participating countries and investigators

Brazil: Cora Luiza Araújo, Cesar G. Victora, Elaine Albernaz, Elaine Tomasi, Rita de Cássia Fossati da Silveira, Gisele Nader (Departamento de Nutrição and Departamento de Medicina Social, Universidade Federal de Pelotas; and Núcleo de Pediatria and Escola de Psicologia, Universidade Católica de Pelotas).

Ghana: Anna Lartey, William B. Owusu, Isabella Sagoe-Moses, Veronica Gomez, Charles Sagoe-Moses (Department of Nutrition and Food Science, University of Ghana; and Ghana Health Service).

India: Nita Bhandari, Maharaj K. Bhan, Sunita Taneja, Temsunaro Rongsen, Jyotsna Chetia, Pooja Sharma, Rajiv Bahl (All India Institute of Medical Sciences).

Norway: Gunn-Elin Aa. Bjoerneboe, Anne Baerug, Elisabeth Tufte, Kaare R. Norum, Karin Rudvin, Hilde Nysaether (Directorate of Health and Social Affairs; National Breastfeeding Centre, Rikshospitalet University Hospital; and Institute for Nutrition Research, University of Oslo).

Oman: Ali Jaffer Mohamed, Deena Alasfoor, Nitya S. Prakash, Ruth M. Mabry, Hanadi Jamaan Al Rajab, Sahar Abdou Helmi (Ministry of Health).

USA: Kathryn G. Dewey, Laurie A. Nommsen-Rivers, Roberta J. Cohen, M. Jane Heinig (University of California, Davis).

Acknowledgements

The WHO Child Growth Standards were constructed by the Coordinating Team in the Department of Nutrition for Health and Development of the World Health Organization.

The Study Group is indebted to the parents, children and more than 200 field staff that participated in the WHO Multicentre Growth Reference Study. The generous contribution of many individuals that provided expertise and advice was also crucial to the development of the growth standards.

The project has received funding from the Bill & Melinda Gates Foundation, the Netherlands Minister for Development Cooperation, the Norwegian Royal Ministry of Foreign Affairs, and the United States Department of Agriculture (USDA). Financial support was also provided by the Ministry of Health of Oman, the United States National Institutes of Health, the Brazilian Ministry of Health and Ministry of Science and Technology, the Canadian International Development Agency, the United Nations University, the Arab Gulf Fund for United Nations Development, the Office of the WHO Representative to India, and the Department of Child and Adolescent Health and Development.

Contents

| | |
|--|------|
| Executive summary | xvii |
| 1. Introduction | 1 |
| 2. Methodology | 3 |
| 2.1 Design of the WHO Multicentre Growth Reference Study | 3 |
| 2.2 Anthropometry methods | 3 |
| 2.3 Sample description | 4 |
| 2.4 Data cleaning procedures and exclusions | 5 |
| 2.5 Statistical methods for constructing the growth curves | 7 |
| 3. Construction of the length/height-for-age standards | 13 |
| 3.1 Indicator-specific methodology | 13 |
| 3.2 Length/height-for-age for boys | 13 |
| 3.2.1 Sample size | 13 |
| 3.2.2 Model selection and results | 14 |
| 3.2.3 WHO standards and their comparison with NCHS and CDC 2000 references | 32 |
| Charts | 33 |
| Tables | 37 |
| Comparison with NCHS | 45 |
| Comparison with CDC 2000 | 46 |
| 3.3 Length/height-for-age for girls | 47 |
| 3.3.1 Sample size | 47 |
| 3.3.2 Model selection and results | 47 |
| 3.3.3 WHO standards and their comparison with NCHS and CDC 2000 references | 59 |
| Charts | 60 |
| Tables | 64 |
| Comparison with NCHS | 72 |
| Comparison with CDC 2000 | 73 |
| 3.4 Comparisons between boys and girls | 74 |
| 3.4.1 WHO | 75 |
| 3.4.2 NCHS | 76 |
| 3.4.3 CDC 2000 | 77 |
| 4. Construction of the weight-for-age standards | 79 |
| 4.1 Indicator-specific methodology | 79 |
| 4.2 Weight-for-age for boys | 79 |
| 4.2.1 Sample size | 79 |
| 4.2.2 Model selection and results | 79 |
| 4.2.3 WHO standards and their comparison with NCHS and CDC 2000 references | 94 |
| Charts | 95 |
| Tables | 97 |
| Comparison with NCHS | 105 |
| Comparison with CDC 2000 | 106 |
| 4.3 Weight-for-age for girls | 107 |
| 4.3.1 Sample size | 107 |
| 4.3.2 Model selection and results | 107 |
| 4.3.3 WHO standards and their comparison with NCHS and CDC 2000 references | 122 |
| Charts | 123 |
| Tables | 125 |
| Comparison with NCHS | 133 |
| Comparison with CDC 2000 | 134 |
| 4.4 Comparisons between boys and girls | 135 |
| 4.4.1 WHO | 136 |
| 4.4.2 NCHS | 137 |

| | | |
|------------------------------------|--|-----|
| 4.4.3 | CDC 2000 | 138 |
| 5. | Construction of the weight-for-length and weight-for-height standards | 139 |
| 5.1 | Indicator-specific methodology | 139 |
| 5.2 | Weight-for-length/height for boys | 139 |
| 5.2.1 | Sample size | 139 |
| 5.2.2 | Model selection and results | 140 |
| 5.2.3 | WHO standards and their comparison with NCHS and CDC 2000 references | 153 |
| Charts | | 154 |
| Tables | | 158 |
| Comparison with NCHS | | 176 |
| Comparison with CDC 2000 | | 178 |
| 5.3 | Weight-for-length/height for girls | 180 |
| 5.3.1 | Sample size | 180 |
| 5.3.2 | Model selection and results | 180 |
| 5.3.3 | WHO standards and their comparison with NCHS and CDC 2000 references | 194 |
| Charts | | 195 |
| Tables | | 199 |
| Comparison with NCHS | | 217 |
| Comparison with CDC 2000 | | 219 |
| 5.4 | Comparisons between boys and girls | 221 |
| 5.4.1 | WHO | 222 |
| 5.4.2 | NCHS | 224 |
| 5.4.3 | CDC 2000 | 226 |
| 6. | Construction of the body mass index-for-age standards | 229 |
| 6.1 | Indicator-specific methodology | 229 |
| 6.2 | BMI-for-age for boys | 230 |
| 6.2.1 | Sample size | 230 |
| 6.2.2 | Model selection and results | 230 |
| Length-based BMI-for-age for boys | | 230 |
| Height-based BMI-for-age for boys | | 241 |
| 6.2.3 | WHO standards and their comparison with CDC 2000 reference | 249 |
| Charts | | 250 |
| Tables | | 254 |
| Comparison with CDC 2000 | | 262 |
| 6.3 | BMI-for-age for girls | 263 |
| 6.3.1 | Sample size | 263 |
| 6.3.2 | Model selection and results | 263 |
| Length-based BMI-for-age for girls | | 263 |
| Height-based BMI-for-age for girls | | 275 |
| 6.3.3 | WHO standards and their comparison with CDC 2000 reference | 284 |
| Charts | | 285 |
| Tables | | 289 |
| Comparison with CDC 2000 | | 297 |
| 6.4 | Comparisons between boys and girls | 298 |
| 6.4.1 | WHO | 299 |
| 6.4.2 | CDC 2000 | 300 |
| 7. | Computation of centiles and z-scores for length/height-for-age, weight-for-age, weight-for-length, weight-for-height and BMI-for-age | 301 |
| 8. | Conclusion | 305 |
| 9. | Bibliography | 309 |
| Appendix A. | Model specifications of the WHO child growth standards | 312 |

Figures

| | | |
|-----------|--|----|
| Figure 1 | Worm plots of z-scores for candidate model with $df(\mu)=11$ and $df(\sigma)=6$ with age transformation $age^{0.35}$ for length/height-for-age for boys | 16 |
| Figure 2 | Fitting of μ and σ curves of Model 1 for length/height-for-age for boys (dotted line) and their respective sample estimates (points with solid line) | 17 |
| Figure 3 | Centile residuals from fitting Model 1 for length/height-for-age from 0 to 24 months for boys | 18 |
| Figure 4 | Centile residuals from fitting Model 1 for length/height-for-age from 24 to 71 months for boys | 18 |
| Figure 5 | Worm plots of z-scores for Model 1 for length/height-for-age for boys | 19 |
| Figure 6 | Worm plots of z-scores for Model 2 for length/height-for-age for boys | 24 |
| Figure 7 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: length-for-age for boys from birth to 24 months | 28 |
| Figure 8 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: length-for-age for boys from birth to 24 months | 29 |
| Figure 9 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: height-for-age for boys from 24 to 71 months | 30 |
| Figure 10 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: height-for-age for boys from 24 to 71 months | 31 |
| Figure 11 | WHO length-for-age z-scores for boys from birth to 24 months | 33 |
| Figure 12 | WHO height-for-age z-scores for boys from 24 to 60 months | 34 |
| Figure 13 | WHO length-for-age percentiles for boys from birth to 24 months | 35 |
| Figure 14 | WHO height-for-age percentiles for boys from 24 to 60 months | 36 |
| Figure 15 | Comparison of WHO with NCHS length/height-for-age z-scores for boys | 45 |
| Figure 16 | Comparison of WHO with CDC 2000 length/height-for-age z-scores for boys | 46 |
| Figure 17 | Fitting of the μ and σ curves of Model 1 for length/height-for-age for girls (dotted line) and their respective sample estimates (points with solid line) | 49 |
| Figure 18 | Centile residuals from fitting Model 1 for length/height-for-age from 0 to 24 months for girls | 50 |
| Figure 19 | Centile residuals from fitting Model 1 for length/height-for-age from 24 to 71 months for girls | 50 |
| Figure 20 | Worm plots of z-scores for Model 1 for length/height-for-age for girls | 51 |
| Figure 21 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: length-for-age for girls from birth to 24 months | 55 |
| Figure 22 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: length-for-age for girls from birth to 24 months | 56 |
| Figure 23 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: height-for-age for girls from 24 to 71 months | 57 |
| Figure 24 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: height-for-age for girls from 24 to 71 months | 58 |
| Figure 25 | WHO length-for-age z-scores for girls from birth to 24 months | 60 |
| Figure 26 | WHO height-for-age z-scores for girls from 24 to 60 months | 61 |

| | | |
|-----------|---|-----|
| Figure 27 | WHO length-for-age percentiles for girls from birth to 24 months | 62 |
| Figure 28 | WHO height-for-age percentiles for girls from 24 to 60 months | 63 |
| Figure 29 | Comparison of WHO with NCHS length/height-for-age z-scores for girls | 72 |
| Figure 30 | Comparison of WHO with CDC 2000 length/height-for-age z-scores for girls | 73 |
| Figure 31 | Comparison of boys' and girls' WHO length/height-for-age z-scores | 75 |
| Figure 32 | Comparison of boys' and girls' NCHS length/height-for-age z-scores | 76 |
| Figure 33 | Comparison of boys' and girls' CDC 2000 length/height-for-age z-scores | 77 |
| Figure 34 | Worm plots of z-scores for Model 1 for weight-for-age for boys | 81 |
| Figure 35 | Fitting of the μ , σ , and ν curves of Model 2 for weight-for-age for boys from 0 to 71 months (dotted line) and their respective sample estimates (points with solid line) | 84 |
| Figure 36 | Centile residuals from fitting Model 2 for weight-for-age from 0 to 24 months for boys | 85 |
| Figure 37 | Centile residuals from fitting Model 2 for weight-for-age from 24 to 71 months for boys | 85 |
| Figure 38 | Worm plots of z-scores for Model 2 for weight-for-age for boys | 86 |
| Figure 39 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: weight-for-age for boys from birth to 24 months | 90 |
| Figure 40 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: weight-for-age for boys from birth to 24 months | 91 |
| Figure 41 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: weight-for-age for boys from 24 to 71 months | 92 |
| Figure 42 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: weight-for-age for boys from 24 to 71 months | 93 |
| Figure 43 | WHO weight-for-age z-scores for boys from birth to 60 months | 95 |
| Figure 44 | WHO weight-for-age percentiles for boys from birth to 60 months | 96 |
| Figure 45 | Comparison of WHO with NCHS weight-for-age z-scores for boys | 105 |
| Figure 46 | Comparison of WHO with CDC 2000 weight-for-age z-scores for boys | 106 |
| Figure 47 | Cubic splines fitted for the ν curve with varying numbers of degrees of freedom | 110 |
| Figure 48 | Fitting of the μ , σ , and ν curves of Model 3 for weight-for-age for girls from 0 to 71 months (dotted line) and their respective sample estimates (points with solid line) | 113 |
| Figure 49 | Centile residuals from fitting Model 3 for weight-for-age from 0 to 24 months for girls | 114 |
| Figure 50 | Centile residuals from fitting Model 3 for weight-for-age from 24 to 71 months for girls | 114 |
| Figure 51 | Worm plots of z-scores for Model 3 for weight-for-age for girls | 115 |
| Figure 52 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: weight-for-age for girls from birth to 24 months | 118 |
| Figure 53 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: weight-for-age for girls from birth to 24 months | 119 |

| | | |
|-----------|---|-----|
| Figure 54 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: weight-for-age for girls from 24 to 71 months | 120 |
| Figure 55 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: weight-for-age for girls from 24 to 71 months | 121 |
| Figure 56 | WHO weight-for-age z-scores for girls from birth to 60 months | 123 |
| Figure 57 | WHO weight-for-age percentiles for girls from birth to 60 months | 124 |
| Figure 58 | Comparison of WHO with NCHS weight-for-age z-scores for girls | 133 |
| Figure 59 | Comparison of WHO with CDC 2000 weight-for-age z-scores for girls | 134 |
| Figure 60 | Comparison of boys' and girls' WHO weight-for-age z-scores | 136 |
| Figure 61 | Comparison of boys' and girls' NCHS weight-for-age z-scores | 137 |
| Figure 62 | Comparison of boys' and girls' CDC 2000 weight-for-age z-scores | 138 |
| Figure 63 | Worm plots of z-scores for Model 1 for weight-for-length/height for boys | 141 |
| Figure 64 | Fitting of the μ , σ , and ν curves of Model 2 for weight-for-length/height for boys (dotted line) and their respective sample estimates (points with solid line) | 143 |
| Figure 65 | Centile residuals from fitting Model 2 for weight-for-length/height for boys | 144 |
| Figure 66 | Worm plots of z-scores for Model 2 for weight-for-length/height for boys | 146 |
| Figure 67 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: weight-for-length for boys | 149 |
| Figure 68 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: weight-for-length for boys | 150 |
| Figure 69 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: weight-for-height for boys | 151 |
| Figure 70 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: weight-for-height for boys | 152 |
| Figure 71 | WHO weight-for-length z-scores for boys from 45 to 110 cm | 154 |
| Figure 72 | WHO weight-for-height z-scores for boys from 65 to 120 cm | 155 |
| Figure 73 | WHO weight-for-length percentiles for boys from 45 to 110 cm | 156 |
| Figure 74 | WHO weight-for-height percentiles for boys from 65 to 120 cm | 157 |
| Figure 75 | Comparison of WHO with NCHS weight-for-length z-scores for boys | 176 |
| Figure 76 | Comparison of WHO with NCHS weight-for-height z-scores for boys | 177 |
| Figure 77 | Comparison of WHO with CDC 2000 weight-for-length z-scores for boys | 178 |
| Figure 78 | Comparison of WHO with CDC 2000 weight-for-height z-scores for boys | 179 |
| Figure 79 | Fitting of the μ , σ , and ν curves of Model 2 for weight-for-length/height for girls (dotted line) and their respective sample estimates (points with solid line) | 183 |
| Figure 80 | Centile residuals from fitting Model 2 for weight-for-length/height for girls | 184 |
| Figure 81 | Worm plots of z-scores for Model 2 for weight-for-length/height for girls | 186 |
| Figure 82 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: weight-for-length for girls | 190 |

| | | |
|------------|---|-----|
| Figure 83 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: weight-for-length for girls | 191 |
| Figure 84 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: weight-for-height for girls | 192 |
| Figure 85 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: weight-for-height for girls | 193 |
| Figure 86 | WHO weight-for-length z-scores for girls | 195 |
| Figure 87 | WHO weight-for-height z-scores for girls | 196 |
| Figure 88 | WHO weight-for-length percentiles for girls | 197 |
| Figure 89 | WHO weight-for-height percentiles for girls | 198 |
| Figure 90 | Comparison of WHO with NCHS weight-for-length z-scores for girls | 217 |
| Figure 91 | Comparison of WHO with NCHS weight-for-height z-scores for girls | 218 |
| Figure 92 | Comparison of WHO with CDC 2000 weight-for-length z-scores for girls | 219 |
| Figure 93 | Comparison of WHO with CDC 2000 weight-for-height z-scores for girls | 220 |
| Figure 94 | Comparison of boys' and girls' WHO weight-for-length z-scores | 222 |
| Figure 95 | Comparison of boys' and girls' WHO weight-for-height z-scores | 223 |
| Figure 96 | Comparison of boys' and girls' NCHS weight-for-length z-scores | 224 |
| Figure 97 | Comparison of boys' and girls' NCHS weight-for-height z-scores | 225 |
| Figure 98 | Comparison of boys' and girls' CDC 2000 weight-for-length z-scores | 226 |
| Figure 99 | Comparison of boys' and girls' CDC 2000 weight-for-height z-scores | 227 |
| Figure 100 | Worm plots of z-scores for Model 1 for length-based BMI-for-age for boys | 232 |
| Figure 101 | Fitting of the μ , σ , and v curves of Model 2 for length-based BMI-for-age from 0 to 24 months for boys (dotted line) and their respective sample estimates (points with solid line) | 234 |
| Figure 102 | Centile residuals from fitting Model 2 for length-based BMI-for-age from 0 to 24 months for boys | 235 |
| Figure 103 | Worm plots of z-scores for Model 2 for length-based BMI-for-age for boys | 236 |
| Figure 104 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: length-based BMI-for-age for boys from birth to 24 months | 239 |
| Figure 105 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: length-based BMI-for-age for boys from birth to 24 months | 240 |
| Figure 106 | Fitting of the μ , σ , and v curves of Model 1 for height-based BMI-for-age from 18 to 71 months for boys (dotted line) and their respective sample estimates (points with solid line) | 242 |
| Figure 107 | Centile residuals from fitting Model 1 for height-based BMI-for-age from 18 to 71 months for boys | 243 |
| Figure 108 | Worm plots of z-scores for Model 1 for height-based BMI-for-age for boys | 246 |
| Figure 109 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: height-based BMI-for-age for boys from 24 to 71 months | 247 |
| Figure 110 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: height-based BMI-for-age for boys from 24 to 71 months | 248 |

| | | |
|------------|--|-----|
| Figure 111 | WHO length-based BMI-for-age z-scores for boys from birth to 24 months | 250 |
| Figure 112 | WHO height-based BMI-for-age z-scores for boys from 24 to 60 months | 251 |
| Figure 113 | WHO length-based BMI-for-age percentiles for boys from birth to 24 months | 252 |
| Figure 114 | WHO height-based BMI-for-age percentiles for boys from 24 to 60 months | 253 |
| Figure 115 | Comparison of WHO with CDC 2000 BMI-for-age z-scores for boys | 262 |
| Figure 116 | Cubic splines fitted for the σ curve with varying numbers of degrees of freedom | 265 |
| Figure 117 | Worm plots of z-scores for Model 1 for length-based BMI-for-age for girls | 267 |
| Figure 118 | Fitting of the μ , σ , and ν curves of Model 2 for length-based BMI-for-age from 0 to 24 months for girls (dotted line) and their respective sample estimates (points with solid line) | 268 |
| Figure 119 | Centile residuals from fitting Model 2 for length-based BMI-for-age from 0 to 24 months for girls | 269 |
| Figure 120 | Worm plots of z-scores for Model 2 for length-based BMI-for-age for girls | 270 |
| Figure 121 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: length-based BMI-for-age for girls from birth to 24 months | 273 |
| Figure 122 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: length-based BMI-for-age for girls from birth to 24 months | 274 |
| Figure 123 | Cubic splines fitted for the μ curve with varying numbers of degrees of freedom | 276 |
| Figure 124 | Fitting of the μ , σ , and ν curves of Model 1 for height-based BMI-for-age from 18 to 71 months for girls (dotted line) and their respective sample estimates (points with solid line) | 277 |
| Figure 125 | Centile residuals from fitting Model 1 for height-based BMI-for-age from 18 to 71 months for girls | 278 |
| Figure 126 | Worm plots of z-scores for Model 1 for height-based BMI-for-age for girls | 279 |
| Figure 127 | 3rd, 10th, 50th, 90th, 97th smoothed centile curves and empirical values: height-based BMI-for-age for girls from 24 to 71 months | 282 |
| Figure 128 | 5th, 25th, 50th, 75th, 95th smoothed centile curves and empirical values: height-based BMI-for-age for girls from 24 to 71 months | 283 |
| Figure 129 | WHO length-based BMI-for-age z-scores for girls from birth to 24 months | 285 |
| Figure 130 | WHO height-based BMI-for-age z-scores for girls from 24 to 60 months | 286 |
| Figure 131 | WHO length-based BMI-for-age percentiles for girls from birth to 24 months | 287 |
| Figure 132 | WHO height-based BMI-for-age percentiles for girls from 24 to 60 months | 288 |
| Figure 133 | Comparison of WHO with CDC 2000 BMI-for-age z-scores for girls | 297 |
| Figure 134 | Comparison of boys' and girls' WHO BMI-for-age z-scores | 299 |
| Figure 135 | Comparison of boys' and girls' CDC 2000 BMI-for-age z-scores | 300 |
| Figure 136 | Examples of children ranked according to the WHO BMI-for-age standards | 303 |

Tables

| | | |
|----------|---|----|
| Table 1 | Total sample and number of compliant children in the longitudinal component | 4 |
| Table 2 | Comparison of mean size at birth for compliant newborns and those that contributed only birth measurements | 5 |
| Table 3 | Total sample of children in the cross-sectional component | 5 |
| Table 4 | Total sample of children in the cross-sectional component by number of visits and total number of records | 5 |
| Table 5 | Number of observations by sex and study component included and excluded on the basis of weight-for-length/height | 6 |
| Table 6 | Number of observations used in the construction of the WHO child growth standards by sex and anthropometric indicator | 7 |
| Table 7 | Interpretation of various patterns in the worm plot | 10 |
| Table 8 | Summary of differences between recumbent length and standing height in a sample of children measured both ways | 13 |
| Table 9 | Longitudinal sample sizes for length/height-for-age for boys | 14 |
| Table 10 | Cross-sectional sample sizes for length/height-for-age for boys | 14 |
| Table 11 | Global deviance (GD) for models within the class $BCPE(x=age^\lambda, df(\mu)=10, df(\sigma)=6, v=1, \tau=2)$ for length/height-for-age for boys | 14 |
| Table 12 | Goodness-of-fit summary for models using the BCPE distribution with fixed $v=1$ and $\tau=2$ for length/height-for-age for boys | 15 |
| Table 13 | Observed proportions of children with measurements below the fitted centiles from Model 1, length/height-for-age for boys | 20 |
| Table 14 | Q-test for z-scores from Model 1 [$BCPE(x=age^{0.35}, df(\mu)=12, df(\sigma)=6, v=1, \tau=2)$] for length/height-for-age for boys | 22 |
| Table 15 | Goodness-of-fit summary for models $BCPE(x=age^{0.35}, df(\mu)=12, df(\sigma)=6, df(v)=?, \tau=2)$ for length/height-for-age for boys | 23 |
| Table 16 | Q-test for z-scores from Model 2 [$BCPE(x=age^{0.35}, df(\mu)=12, df(\sigma)=6, df(v)=6, \tau=2)$] for length/height-for-age for boys | 25 |
| Table 17 | Observed proportions of children with measurements below the fitted centiles from Model 2, length/height-for-age for boys | 26 |
| Table 18 | Length-for-age for boys, age in weeks | 37 |
| Table 19 | Length-for-age for boys, age in years and months | 39 |
| Table 20 | Height-for age for boys, age in years and months | 41 |
| Table 21 | Longitudinal sample sizes for length/height-for-age for girls | 47 |
| Table 22 | Cross-sectional sample sizes for length/height-for-age for girls | 47 |
| Table 23 | Global deviance (GD) for models within the class $BCPE(x=age^\lambda, df(\mu)=10, df(\sigma)=6, v=1, \tau=2)$ for length/height-for-age for girls | 48 |
| Table 24 | Goodness-of-fit summary for models using the BCPE distribution with fixed $v=1$ and $\tau=2$ for length/height-for-age for girls | 48 |
| Table 25 | Observed proportions of children with measurements below the fitted centiles from Model 1, length/height-for-age for girls | 52 |

| | | |
|----------|---|-----|
| Table 26 | Q-test for z-scores from Model 1 [BCPE($x=age^{0.35}$, $df(\mu)=10$, $df(\sigma)=5$, $v=1$, $\tau=2$)] for length/height-for-age for girls | 54 |
| Table 27 | Length-for-age for girls, age in weeks | 64 |
| Table 28 | Length-for-age for girls, age in years and months | 66 |
| Table 29 | Height-for-age for girls, age in years and months | 68 |
| Table 30 | Longitudinal sample sizes for weight-for-age for boys | 79 |
| Table 31 | Cross-sectional sample sizes for weight-for-age for boys | 79 |
| Table 32 | Global deviance (GD) for models within the class BCPE($x=age^\lambda$, $df(\mu)=9$, $df(\sigma)=4$, $df(v)=4$, $\tau=2$) for weight-for-age for boys | 80 |
| Table 33 | Goodness-of-fit summary for models using the BCPE distribution with fixed $v=1$ and $\tau=2$ for weight-for-age for boys | 80 |
| Table 34 | Q-test for z-scores from Model 1 [BCPE($x=age^{0.35}$, $df(\mu)=11$, $df(\sigma)=7$, $v=1$, $\tau=2$)] for weight-for-age for boys | 82 |
| Table 35 | Goodness-of-fit summary for models BCPE($x=age^{0.35}$, $df(\mu)=11$, $df(\sigma)=7$, $df(v)=?$, $\tau=2$) for weight-for-age for boys | 83 |
| Table 36 | Q-test for z-scores from Model 2 [BCPE($x=age^{0.35}$, $df(\mu)=11$, $df(\sigma)=7$, $df(v)=2$, $\tau=2$)] for weight-for-age for boys | 87 |
| Table 37 | Observed proportions of children with measurements below the fitted centiles from Model 2, weight-for-age for boys | 88 |
| Table 38 | Weight-for-age for boys, age in weeks | 97 |
| Table 39 | Weight-for-age for boys, age in years and months | 99 |
| Table 40 | Longitudinal sample sizes for weight-for-age for girls | 107 |
| Table 41 | Cross-sectional sample sizes for weight-for-age for girls | 107 |
| Table 42 | Global deviance (GD) for models within the class BCPE($x=age^\lambda$, $df(\mu)=9$, $df(\sigma)=4$, $df(v)=4$, $\tau=2$) for weight-for-age for girls | 108 |
| Table 43 | Goodness-of-fit summary for models using the BCPE distribution with fixed $v=1$ and $\tau=2$ for weight-for-age for girls | 108 |
| Table 44 | Q-test for z-scores from Model 1 [BCPE($x=age^{0.35}$, $df(\mu)=11$, $df(\sigma)=7$, $v=1$, $\tau=2$)] for weight-for-age for girls | 109 |
| Table 45 | Goodness-of-fit summary for models BCPE($x=age^{0.35}$, $df(\mu)=11$, $df(\sigma)=7$, $df(v)=?$, $\tau=2$) for weight-for-age for girls | 110 |
| Table 46 | Q-test for z-scores from Model 2 [BCPE($x=age^{0.35}$, $df(\mu)=11$, $df(\sigma)=7$, $df(v)=5$, $\tau=2$)] for weight-for-age for girls | 111 |
| Table 47 | Q-test for z-scores from Model 3 [BCPE($x=age^{0.35}$, $df(\mu)=11$, $df(\sigma)=7$, $df(v)=3$, $\tau=2$)] for weight-for-age for girls | 112 |
| Table 48 | Observed proportions of children with measurements below the fitted centiles from Model 3, weight-for-age for girls | 116 |
| Table 49 | Weight-for-age for girls, age in weeks | 125 |
| Table 50 | Weight-for-age for girls, age in years and months | 127 |
| Table 51 | Sample sizes for boys' weight-for-length/height by length interval | 139 |
| Table 52 | Goodness-of-fit summary for models using the BCPE distribution with fixed $v=1$ and $\tau=2$ for weight-for-length/height for boys | 140 |

| | | |
|----------|---|-----|
| Table 53 | Q-test for z-scores from Model 1 [BCPE($x=length$ (or $height+0.7$), $df(\mu)=13$, $df(\sigma)=6$, $v=1$, $\tau=2$)] for weight-for-length/height for boys | 142 |
| Table 54 | Goodness-of-fit summary for models BCPE($x=length$ (or $height+0.7$), $df(\mu)=13$, $df(\sigma)=6$, $df(v)=?$, $\tau=2$) for weight-for-length/height for boys | 142 |
| Table 55 | Q-test for z-scores from Model 2 [BCPE($x=length$ (or $height+0.7$), $df(\mu)=13$, $df(\sigma)=6$, $df(v)=1$, $\tau=2$)] for weight-for-length/height for boys | 145 |
| Table 56 | Observed proportions of children with measurements below the fitted centiles from Model 2, weight-for-length/height for boys | 147 |
| Table 57 | Weight-for-length for boys | 158 |
| Table 58 | Weight-for-height for boys | 168 |
| Table 59 | Sample sizes for girls' weight-for-length/height by length interval | 180 |
| Table 60 | Goodness-of-fit summary for models using the BCPE distribution with fixed $v=1$ and $\tau=2$ for weight-for-length/height for girls | 181 |
| Table 61 | Q-test for z-scores from Model 1 [BCPE($x=length$ (or $height+0.7$), $df(\mu)=12$, $df(\sigma)=4$, $v=1$, $\tau=2$)] for weight-for-length/height for girls | 182 |
| Table 62 | Goodness-of-fit summary for models BCPE($x=length$ (or $height+0.7$), $df(\mu)=12$, $df(\sigma)=4$, $df(v)=?$, $\tau=2$) for weight-for-length/height for girls | 183 |
| Table 63 | Q-test for z-scores from Model 2 [BCPE($x=length$ (or $height+0.7$), $df(\mu)=12$, $df(\sigma)=4$, $df(v)=1$, $\tau=2$)] for weight-for-length/height for girls | 185 |
| Table 64 | Q-test for z-scores from model BCPE($x=length$ (or $height+0.7$), $df(\mu)=12$, $df(\sigma)=4$, $df(v)=1$, $\tau=2.13$) for weight-for-length/height for girls | 187 |
| Table 65 | Observed proportions of children with measurements below the fitted centiles from Model 2, weight-for-length/height for girls | 188 |
| Table 66 | Weight-for-length for girls | 199 |
| Table 67 | Weight-for-height for girls | 209 |
| Table 68 | Longitudinal sample sizes for BMI-for-age for boys | 230 |
| Table 69 | Cross-sectional sample sizes for BMI-for-age for boys | 230 |
| Table 70 | Global deviance (GD) for models within the class BCPE($x=age^\lambda$, $df(\mu)=9$, $df(\sigma)=4$, $df(v)=4$, $\tau=2$) for length-based BMI-for-age for boys | 231 |
| Table 71 | Goodness-of-fit summary for models using the BCPE distribution with fixed $v=1$ and $\tau=2$ for length-based BMI-for-age for boys | 231 |
| Table 72 | Q-test for z-scores from Model 1 [BCPE($x=age^{0.05}$, $df(\mu)=10$, $df(\sigma)=4$, $v=1$, $\tau=2$)] for length-based BMI-for-age for boys | 233 |
| Table 73 | Goodness-of-fit summary for models BCPE($x=age^{0.05}$, $df(\mu)=10$, $df(\sigma)=4$, $df(v)=?$, $\tau=2$) for length-based BMI-for-age for boys | 233 |
| Table 74 | Q-test for z-scores from Model 2 [BCPE($x=age^{0.05}$, $df(\mu)=10$, $df(\sigma)=4$, $df(v)=3$, $\tau=2$)] for length-based BMI-for-age for boys | 237 |
| Table 75 | Observed proportions of children with measurements below the fitted centiles from Model 2, length-based BMI-for-age for boys | 238 |
| Table 76 | Goodness-of-fit summary for models using the BCPE distribution with fixed $v=1$ and $\tau=2$ for height-based BMI-for-age for boys | 241 |
| Table 77 | Goodness-of-fit summary for models BCPE($x=age$, $df(\mu)=4$, $df(\sigma)=3$, $df(v)=?$, $\tau=2$) for height-based BMI-for-age for boys | 242 |

| | | |
|----------|---|-----|
| Table 78 | Observed proportions of children with measurements below the fitted centiles from Model 1, height-based BMI-for-age for boys | 244 |
| Table 79 | Q-test for z-scores from Model 1 [BCPE($x=age$, $df(\mu)=4$, $df(\sigma)=3$, $df(v)=3$, $\tau=2$)] for height-based BMI-for-age for boys | 245 |
| Table 80 | Length-based BMI-for-age for boys, age in weeks | 254 |
| Table 81 | Length-based BMI-for-age for boys, age in years and months | 256 |
| Table 82 | Height-based BMI-for-age for boys, age in years and months | 258 |
| Table 83 | Longitudinal sample sizes for BMI-for-age for girls | 263 |
| Table 84 | Cross-sectional sample sizes for BMI-for-age for girls | 263 |
| Table 85 | Global deviance (GD) for models within the class BCPE($x=age^\lambda$, $df(\mu)=9$, $df(\sigma)=4$, $df(v)=4$, $\tau=2$) for length-based BMI-for-age for girls | 264 |
| Table 86 | Goodness-of-fit summary for models using the BCPE distribution with fixed $v=1$ and $\tau=2$ for length-based BMI-for-age for girls | 264 |
| Table 87 | Q-test for z-scores from Model 1 [BCPE($x=age^{0.05}$, $df(\mu)=10$, $df(\sigma)=3$, $v=1$, $\tau=2$)] for length-based BMI-for-age for girls | 266 |
| Table 88 | Goodness-of-fit summary for models BCPE($x=age^{0.05}$, $df(\mu)=10$, $df(\sigma)=3$, $df(v)=?$, $\tau=2$) for length-based BMI-for-age for girls | 267 |
| Table 89 | Q-test for z-scores from Model 2 [BCPE($x=age^{0.05}$, $df(\mu)=10$, $df(\sigma)=3$, $df(v)=3$, $\tau=2$)] for length-based BMI-for-age for girls | 271 |
| Table 90 | Observed proportions of children with measurements below the fitted centiles from Model 2 for length-based BMI-for-age for girls | 272 |
| Table 91 | Goodness-of-fit summary for models using the BCPE distribution with fixed $v=1$ and $\tau=2$ for height-based BMI-for-age for girls | 275 |
| Table 92 | Goodness-of-fit summary for models BCPE($x=age$, $df(\mu)=4$, $df(\sigma)=4$, $df(v)=?$, $\tau=2$) for height-based BMI-for-age for girls | 276 |
| Table 93 | Q-test for z-scores from Model 1 [BCPE($x=age$, $df(\mu)=4$, $df(\sigma)=4$, $df(v)=1$, $\tau=2$)] for height-based BMI-for age for girls | 280 |
| Table 94 | Observed proportions of children with measurements below the fitted centiles from Model 1 for height-based BMI-for-age for girls | 281 |
| Table 95 | Length-based BMI-for-age for girls, age in weeks | 289 |
| Table 96 | Length-based BMI-for-age for girls, age in years and months | 291 |
| Table 97 | Height-based BMI-for-age for girls, age in years and months | 293 |