



# ESSENTIAL INTERVENTIONS, COMMODITIES AND GUIDELINES

for Reproductive,  
Maternal, Newborn  
and Child Health



**A GLOBAL REVIEW OF THE KEY INTERVENTIONS  
RELATED TO REPRODUCTIVE, MATERNAL, NEWBORN  
AND CHILD HEALTH (RMNCH)**

This document is designed for an audience of policy-makers who seek information on the specific health interventions to address the main causes of maternal, newborn and child deaths.

It is the result of collaborative work among many partners. The process was led by the World Health Organization, Switzerland, and the Aga Khan University, Pakistan. Experts in maternal, newborn and child health participated in meetings in Geneva in April 2010 and September 2011 and provided inputs to the development and finalization of this document. The contributions of the World Health Organization, the Aga Khan University, invited experts and partners are gratefully acknowledged.

This publication, and related advocacy material, will be distributed to over 430 PMNCH partners, and other stakeholders, primarily via the PMNCH website and knowledge portal. In addition, it will be distributed, and discussed, at selected RMNCH advocacy events.

Publication reference: The Partnership for Maternal, Newborn & Child Health. 2011. *A Global Review of the Key Interventions Related to Reproductive, Maternal, Newborn and Child Health (RMNCH)*. Geneva, Switzerland: PMNCH.

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

**Photos:** Front cover, © UNICEF/NYHQ2006-0779/Shehzad Noorani, UN Photo/Eskinder Debebe and UNICEF/BANA2006-01117/Munira Munni; page 16, WHO/Christopher Black; page 18, Joshua Roberts/Save the Children (Mali); page 21, UN Photo/Albert Gonzalez Farran; page 22, UNICEF/060990G; back cover, WHO/Christopher Black, UNICEF/Shehzad Noorani, UNI58124 and UN Photo/Evan Schneider.

**Design:** Roberta Annovi.

<b>ONE-PAGE SUMMARY OF ESSENTIAL INTERVENTIONS</b>	Pg. 4
<b>EXECUTIVE SUMMARY</b>	Pg. 6
Why reproductive, maternal, newborn and child health?	Pg. 6
Methodology	Pg. 7
Evidence-based findings	Pg. 10
<b>REPRODUCTIVE AND MATERNAL HEALTH INTERVENTIONS</b>	Pg. 12
<b>NEWBORN CARE INTERVENTIONS</b>	Pg. 17
<b>CHILD HEALTH INTERVENTIONS</b>	Pg. 19
<b>CROSS-CUTTING COMMUNITY STRATEGIES</b>	Pg. 22
<b>BIBLIOGRAPHY</b>	Pg. 23
<b>ACKNOWLEDGEMENTS</b>	Pg. 26

# ONE-PAGE SUMMARY OF ESSENTIAL INTERVENTIONS

## ESSENTIAL, EVIDENCE-BASED INTERVENTIONS TO REDUCE REPRODUCTIVE, MATERNAL,

CONTINUUM OF CARE	ADOLESCENCE & PRE-PREGNANCY	PREGNANCY (ANTENATAL)	CHILDBIRTH
<b>ALL LEVELS:</b>  <b>COMMUNITY PRIMARY REFERRAL</b>	<ul style="list-style-type: none"> <li>Family planning (advice, hormonal and barrier methods)</li> <li>Prevent and manage sexually transmitted infections, HIV</li> <li>Folic acid fortification/supplementation to prevent neural tube defects</li> </ul>	<ul style="list-style-type: none"> <li>Iron and folic acid supplementation</li> <li>Tetanus vaccination</li> <li>Prevention and management of malaria with insecticide treated nets and antimalarial medicines</li> <li>Prevention and management of sexually transmitted infections and HIV, including with antiretroviral medicines</li> <li>Calcium supplementation to prevent hypertension (high blood pressure)</li> <li>Interventions for cessation of smoking</li> </ul>	<ul style="list-style-type: none"> <li>Prophylactic uterotonics to prevent postpartum haemorrhage (excessive bleeding after birth)</li> <li>Manage postpartum haemorrhage using uterine massage and uterotonics</li> <li>Social support during childbirth</li> </ul>
<b>PRIMARY AND REFERRAL</b>	<ul style="list-style-type: none"> <li>Family planning (hormonal, barrier and selected surgical methods)</li> </ul>	<ul style="list-style-type: none"> <li>Screening for and treatment of syphilis</li> <li>Low dose aspirin to prevent pre-eclampsia</li> <li>Antihypertensive drugs (to treat high blood pressure)</li> <li>Magnesium sulphate for eclampsia</li> <li>Antibiotics for preterm prelabour rupture of membranes</li> <li>Corticosteroids to prevent respiratory distress syndrome in preterm babies</li> <li>Safe abortion</li> <li>Post abortion care</li> </ul>	<ul style="list-style-type: none"> <li>Active management of third stage of labour (to deliver the placenta) to prevent postpartum haemorrhage (<i>as above plus controlled cord traction</i>)</li> <li>Management of postpartum haemorrhage (<i>as above plus manual removal of placenta</i>)</li> <li>Screen and manage HIV (if not already tested)</li> </ul>
<b>REFERRAL*</b>	<ul style="list-style-type: none"> <li>Family planning (surgical methods)</li> </ul>	<ul style="list-style-type: none"> <li>Reduce malpresentation at term with External Cephalic Version</li> <li>Induction of labour to manage prelabour rupture of membranes at term (initiate labour)</li> </ul>	<ul style="list-style-type: none"> <li>Caesarean section for maternal/foetal indication (to save the life of the mother/baby)</li> <li>Prophylactic antibiotic for caesarean section</li> <li>Induction of labour for prolonged pregnancy (initiate labour)</li> <li>Management of postpartum haemorrhage (<i>as above plus surgical procedures</i>)</li> </ul>
<b>COMMUNITY STRATEGIES</b>	<ul style="list-style-type: none"> <li>Home visits for women and children across the continuum of care</li> <li>Women's groups</li> </ul>		

## NEWBORN AND CHILD MORTALITY, AND PROMOTE REPRODUCTIVE HEALTH

POSTNATAL (MOTHER)	POSTNATAL (NEWBORN)	INFANCY & CHILDHOOD
<ul style="list-style-type: none"> <li>Family planning advice and contraceptives</li> <li>Nutrition counselling</li> </ul>	<ul style="list-style-type: none"> <li>Immediate thermal care (to keep the baby warm)</li> <li>Initiation of early breastfeeding (within the first hour)</li> <li>Hygienic cord and skin care</li> </ul>	<ul style="list-style-type: none"> <li>Exclusive breastfeeding for 6 months</li> <li>Continued breastfeeding and complementary feeding from 6 months</li> <li>Prevention and case management of childhood malaria</li> <li>Vitamin A supplementation from 6 months of age</li> <li>Routine immunization plus <i>H.influenzae</i>, meningococcal, pneumococcal and rotavirus vaccines</li> <li>Management of severe acute malnutrition</li> <li>Case management of childhood pneumonia</li> <li>Case management of diarrhoea</li> </ul>
<ul style="list-style-type: none"> <li>Screen for and initiate or continue antiretroviral therapy for HIV</li> <li>Treat maternal anaemia</li> </ul>	<ul style="list-style-type: none"> <li>Neonatal resuscitation with bag and mask (by professional health workers for babies who do not breathe at birth)</li> <li>Kangaroo mother care for preterm (premature) and for less than 2000g babies</li> <li>Extra support for feeding small and preterm babies</li> <li>Management of newborns with jaundice (“yellow” newborns)</li> <li>Initiate prophylactic antiretroviral therapy for babies exposed to HIV</li> </ul>	<ul style="list-style-type: none"> <li>Comprehensive care of children infected with, or exposed to, HIV</li> </ul>
<ul style="list-style-type: none"> <li>Detect and manage postpartum sepsis (serious infections after birth)</li> </ul>	<ul style="list-style-type: none"> <li>Presumptive antibiotic therapy for newborns at risk of bacterial infection</li> <li>Use of surfactant (respiratory medication) to prevent respiratory distress syndrome in preterm babies</li> <li>Continuous positive airway pressure (CPAP) to manage babies with respiratory distress syndrome</li> <li>Case management of neonatal sepsis, meningitis and pneumonia</li> </ul>	<ul style="list-style-type: none"> <li>Case management of meningitis</li> </ul>
<p>* Family planning interventions at Referral level include those provided at the Primary level</p>		

### WHY REPRODUCTIVE, MATERNAL, NEWBORN AND CHILD HEALTH?

Poor maternal, newborn and child health remains a significant problem in developing countries. Worldwide, 358 000 women die during pregnancy and childbirth every year<sup>1</sup> and an estimated 7.6 million children die under the age of five.<sup>2</sup> The majority of maternal deaths occur during or immediately after childbirth. The common medical causes for maternal death include bleeding, high blood pressure, prolonged and obstructed labour, infections and unsafe abortions. A child's risk of dying is highest during the first 28 days of life when about 40% of under-five deaths take place, translating into three million deaths.<sup>2</sup> Up to one half of all newborn deaths occur within the first 24 hours of life and 75% occur in the first week. Globally, the main causes of neonatal death are preterm birth, severe infections and asphyxia. Children in low-income countries are nearly 18 times more likely to die before the age of five than children in high-income countries.<sup>2</sup>

Good maternal health and nutrition are important contributors to child survival. The lack of essential interventions to address these and other health conditions often contribute to indices of neonatal morbidity and mortality (including stillbirths, neonatal deaths and other adverse clinical outcomes).

The highest maternal, neonatal and under-five mortality rates are in sub-Saharan Africa and in Southern Asia.<sup>2</sup> Although substantial progress has been made towards achieving the Millennium Development Goals (MDGs) 4 and 5, the rates of decline in maternal, newborn and under-five mortality remain insufficient to achieve these goals by 2015. Interventions and strategies for improving reproductive, maternal, newborn and child health and survival are closely related and must be provided through a continuum of care approach. When linked together and included as integrated programmes, these interventions can lower costs, promote greater efficiencies and reduce duplication of resources. However, few efforts have been made to identify synergies and integrate these interventions across the continuum of care. Despite the existing plethora of knowledge, there is a lack of consensus on how best to move forward in a coordinated manner so as to achieve progress towards the MDGs. Furthermore, consensus is also needed on the level of evidence.



The foremost aim of this global review is to compile existing evidence on the impact of different interventions on the main causes of maternal, newborn and child deaths. The specific objectives of this review were to serve as a first step towards:

- Developing consensus on the content of RMNCH packages of interventions at each level of the health system across the continuum of care.
- Facilitating the scaling-up of these interventions.
- Identifying research gaps in the content of core packages of interventions.

### *Policy and regulatory environment*

Policy and regulations are crucial to the implementation of any interventions. The recommended list of interventions should be reviewed in light of the existing national policy and regulatory environment. All interventions provided should comply with the laws and policies of the country. When required, these laws and policies may be reviewed and updated to ensure that priority life saving interventions are delivered.



**FIGURE 1:****MDG 4: TRENDS IN UNDER-FIVE MORTALITY, 1990 - 2007****MDG 5: TRENDS IN MATERNAL MORTALITY RATIOS, 2008****METHODOLOGY****Search strategy**

A total of 142 RMNCH interventions were identified, assessed and selected for this review (there is a 700 page compilation of the evidence which underpins this short summary available at the PMNCH),<sup>3</sup> based on current WHO recommendations contained in the following publications: Guidelines on HIV and Infant Feeding (2010); Integrated Management of Childhood Illness (2008); Integrated Management of Childhood Illness for high HIV settings (2008); the Pocket Book on Hospital Care for Children (2005); Integrated Management of Pregnancy and Childbirth Clinical Guidelines (2007); Recommended Interventions for Improving Maternal and Newborn Health - Integrated Management of Pregnancy and Childbirth (2007). Interventions published in the Child and Neonatal Lancet Series (2003 and 2005, respectively) as well as in the WHO Recommended Interventions for Improving Maternal and Newborn health (2010).

Inclusion criteria comprised the following: (i) the intervention has an alleged impact on reducing maternal, neonatal and child mortality; (ii) the intervention is suitable for delivery in low- and middle-income countries, and/or settings where minimal essential care is generally available; and (iii) the intervention is delivered through the health sector (community level up to the referral level of health care).

Relevant reviews for each intervention were identified from the following electronic databases: the Cochrane database of systematic reviews, the Cochrane database of abstract reviews of effectiveness (DARE), the Cochrane database of systematic reviews of randomized control trials (RCTs), and PubMed. The reference lists of reviews and recommendations from experts in the field were also used as sources to obtain additional publications. The principal focus was on the existing systematic reviews and meta-analysis.

### *Selection on interventions*

The interventions were prioritized according to the following criteria:

- Interventions expected to have a **significant impact on maternal, newborn and child survival**, addressing the main causes of maternal, newborn and child mortality.
- Interventions suitable for implementation in **low- and middle-income countries**; minimal essential care.
- Interventions delivered through the **health sector**, from the community up to the first referral level of health service provision.

### *Classification of interventions*

The interventions were classified into categories A, B and C, according to the framework provided in **Box 1**.

#### **Box 1:**

CATEGORY	EVIDENCE FOR INTERVENTION CATEGORIES	DELIVERY STRATEGIES	ACTION
A	Intervention evidence agreed	Delivery strategy agreed	Disseminate for rapid scale-up
B	Intervention evidence agreed	Delivery strategy no consensus	Collate evidence and define gaps in evidence for delivery strategies – seek consensus
C	Intervention evidence still questioned	Delivery strategy no consensus	Further research required

The classification of the effect of interventions according to the evidence available was done based on that used by the Cochrane group, as follows:

A	B	C	D	E
Interventions that are beneficial	Interventions likely to be beneficial	Interventions with a trade-off between beneficial and adverse effects	Interventions of unknown effect, including absence of reviews	Interventions likely to be ineffective or harmful

This classification benefited from being broadly known, recognized and accepted since it is the classification used by the Cochrane systematic review process that has guided this exercise from the beginning. The “evidence” was restricted to published systematic reviews; not including single studies.



The origin of evidence included the following three different levels of delivery of interventions and these were defined in the publication by the World Bank “Priorities in Health”:<sup>4</sup>

### COMMUNITY LEVEL/HOME



**(1) COMMUNITY LEVEL/HOME** - Health care providers at this level include community health workers and outreach workers. It utilizes resources such as volunteers’ time, local knowledge and community confidence and trust as channels for delivery of interventions generally related to safe motherhood, nutrition and simple prevention and treatments. Many countries have attempted to construct links between community-based health care resources and

households for a range of health programmes. These programmes do not substitute for a health system, but provide a channel for reaching families with information and resources. Community health workers (CHWs) not only promote healthy behaviours and preventive action but can mobilize demand for appropriate services at other levels. The success of community health efforts depends critically on the context, including level of development of infrastructure, services and socioeconomic resources.

### FIRST LEVEL/OUTREACH



**(2) FIRST LEVEL/OUTREACH** - Health care providers at this level of care include professionals, outreach workers as well as the community health workers. It includes a range of initiatives that are associated with the Alma Ata Declaration on Primary Health Care approved by WHO in 1978. More recently, the WHO Commission on Macroeconomics and Health described the need for developing services that are close to the client. The basic notion is a common one:

recognition that a certain range of health care services must act as an interface between families and community programmes on the one hand, and hospitals and national health policies on the other. There has been substantial convergence in the content of general first level primary care over time: maternity related care (for instance, prenatal care, skilled birth attendance and family planning), interventions to address childhood diseases (such as vaccine preventable diseases, acute respiratory infections, diarrhoea) and prevention and treatment of major infectious diseases.

### REFERRAL LEVEL/DISTRICT HOSPITAL



**(3) REFERRAL LEVEL** - This level of delivery of interventions refers to hospitals in general. These can be either district hospitals or referral hospitals. The health care providers at this level are professionals.

*District hospitals* - Generally designed to serve people with services that are more sophisticated, technically demanding and specialized than those available at a primary care facility/first level care, but not as specialized

as those provided by referral hospitals. Their range of services includes diagnostics, treatment, care, counselling and rehabilitation. District hospitals may also provide health information, training and administrative and logistical support to primary and community health care programmes. They concentrate skills and resources in one place for the delivery of interventions for conditions that are either uncommon or difficult to treat. They are also a repository of knowledge and diagnostic tools for assessing whether referral to an even more specialized facility is indicated.

**Referral hospitals** - Referral hospitals provide complex clinical care interventions to patients referred from the community, primary/first, or district hospital levels. Referral hospitals need to provide many forms of support, including advice on which patients to refer, proper post discharge care and long-term management of chronic conditions. Referral hospitals can also provide important managerial and administrative support to other facilities, serving as gateways for drugs and medical supplies, laboratory testing services, general procurement, data collection from health information systems and epidemiological surveillance. They are also the vehicle for disseminating technologies by training new staff and providing continuing professional education for existing staff at different facilities.

## EVIDENCE-BASED FINDINGS

The following table lists the interventions classified as “A” based on the criteria defined in Box 1.

### CLASSIFICATION OF INTERVENTIONS ACCORDING TO THE LEVEL OF HEALTH CARE DELIVERY

Intervention	Referral level	1 <sup>st</sup> level	Community
<b>Adolescents &amp; Pre-Pregnancy</b>			
Family planning	✓	✓	✓
Prevent and manage Sexually Transmitted illnesses including Mother-to-Child Transmission of HIV and syphilis	✓	✓	✓
Folic acid fortification and/or supplementation for preventing Neural Tube Defects	✓	✓	✓
<b>Pregnancy</b>			
Management of unintended pregnancy			
▪ Availability and provision of safe abortion care when indicated	✓	-	-
▪ Provision of post abortion care	✓	✓	-
<b>Appropriate antenatal care package:</b>	✓	✓	-
▪ Screening for maternal illnesses			
▪ Screening for hypertensive disorders of pregnancy			
▪ Screening for anaemia			
▪ Iron and folic acid to prevent maternal anaemia			
▪ Tetanus immunization			
▪ Counselling on family planning, birth and emergency preparedness			
▪ Prevention and management of HIV, including with antiretrovirals			
▪ Prevent and manage malaria with insecticide treated nets and antimalarial medicine			
▪ Smoking cessation			
Reduce malpresentation at term with External Cephalic Version	✓	-	-
Prevention of pre-eclampsia			
▪ Calcium to prevent hypertension	✓	✓	-
▪ Low dose aspirin to prevent hypertension	✓	-	-
Magnesium Sulphate for eclampsia	✓	✓	-
Induction of labour to manage prelabour rupture of membranes at term	✓	-	-
Antibiotics for preterm prelabour rupture of membranes	✓	✓	-
Corticosteroids to prevent respiratory distress syndrome in newborns	✓	-	-

Intervention	Referral level	1 <sup>st</sup> level	Community
<b>Childbirth</b>			
Induction of labour for prolonged pregnancy	✓	-	-
Prophylactic uterotonics to prevent postpartum haemorrhage	✓	✓	✓
Active management of third stage of labour to prevent postpartum haemorrhage	✓	✓	-
Management of postpartum haemorrhage (e.g. uterotonics, uterine massage)	✓	✓	✓
Caesarean section for maternal/foetal indication	✓	-	-
Prophylactic antibiotics for caesarean section	✓	-	-
<b>Postnatal (mother)</b>			
Family planning	✓	✓	✓
Prevent and treat maternal anaemia	✓	✓	-
Detect and manage postpartum sepsis	✓	✓	-
Screen and initiate or continue antiretroviral therapy for HIV	✓	✓	-
<b>Postnatal (newborn)</b>			
Immediate thermal care	✓	✓	✓
Initiation of exclusive breastfeeding (within first hour)	✓	✓	✓
Hygienic cord and skin care	✓	✓	✓
Neonatal resuscitation with bag and mask (professional health worker)	✓	✓	-
Case management of neonatal sepsis, meningitis and pneumonia	✓	✓	-
Kangaroo mother care for preterm and for less than 2000g babies	✓	✓	-
Management of newborns with jaundice	✓	✓	-
Surfactant to prevent respiratory distress syndrome in preterm babies	✓	-	-
Continuous positive airway pressure (CPAP) to manage babies with respiratory distress syndrome	✓	-	-
Extra support for feeding small and preterm babies	✓	✓	-
Presumptive antibiotic therapy for newborns at risk of bacterial infections	✓	-	-
<b>Infancy and Childhood</b>			
Exclusive breastfeeding for 6 months	✓	✓	✓
Continued breastfeeding and complementary feeding from 6 months	✓	✓	✓
Prevention and case management of childhood malaria	✓	✓	✓
Vitamin A supplementation from 6 months of age	✓	✓	✓
Comprehensive care of children infected with or exposed to HIV	✓	✓	-
Routine immunization and <i>H. influenzae</i> , meningococcal, pneumococcal and rotavirus vaccines	✓	✓	✓
Management of severe acute malnutrition	✓	✓	-
Case management of childhood pneumonia	✓	✓	✓
Case management of diarrhoea	✓	✓	✓
<b>Cross-cutting community strategies</b>			
Home visits for women and children across the continuum of care	-	-	✓

# REPRODUCTIVE AND MATERNAL HEALTH INTERVENTIONS

PRIORITY INTERVENTIONS	LEVEL OF CARE (COMMUNITY, PRIMARY, REFERRAL)	COMMUNITY OR PROFESSIONAL HEALTH WORKERS	KEY COMMODITIES (SUPPLEMENTED BY ANNEX)	PRACTICE GUIDELINES AND TRAINING MANUALS
<b>PRECONCEPTION/PERICONCEPTUAL INTERVENTIONS</b>				
Family planning <sup>5-7</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Barrier methods (male and female condoms, diaphragm, gels, foams)</li> <li>Oral contraceptives (progestin only and combined)</li> <li>Emergency contraceptives and hormonal injections</li> </ul>	<ul style="list-style-type: none"> <li>Medical eligibility criteria for contraceptive use <a href="http://whqlibdoc.who.int/publications/2010/9789241563888_eng.pdf">http://whqlibdoc.who.int/publications/2010/9789241563888_eng.pdf</a></li> <li>Family Planning: a global handbook for providers <a href="http://whqlibdoc.who.int/publications/2011/9780978856373_eng.pdf">http://whqlibdoc.who.int/publications/2011/9780978856373_eng.pdf</a></li> <li>Surgical Care at the District Hospital <a href="http://www.who.int/surgery/publications/scdh_manual/en/index.html">www.who.int/surgery/publications/scdh_manual/en/index.html</a> pgs 9-8, 11-19</li> </ul>
	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>All of the above plus implants</li> <li>Long acting reversible contraceptives (implants)</li> <li>Intrauterine devices</li> <li>Surgical contraception</li> </ul>	
Prevention and management of Sexually Transmitted Infections (STIs), including HIV for Prevention of Mother-to-Child Transmission (PMTCT) of HIV and syphilis <sup>8,9</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Materials for counselling</li> <li>Condoms (male and female)</li> <li>Antibiotics in line with essential medicine guidelines</li> </ul>	<ul style="list-style-type: none"> <li>Sexually transmitted and other reproductive tract infections: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2005/9241592656.pdf">http://whqlibdoc.who.int/publications/2005/9241592656.pdf</a></li> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> <li>Rapid advice: use of antiretroviral drugs for treating pregnant women and preventing HIV infection in infants <a href="http://www.who.int/hiv/pub/mtct/advice/en/index.html">www.who.int/hiv/pub/mtct/advice/en/index.html</a></li> </ul>
	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Materials for counselling</li> <li>Condoms (male and female)</li> <li>Antibiotics in line with essential medicine guidelines</li> <li>Laboratory test kits for STI/HIV</li> <li>Antiretroviral medicines (refer to the essential list of medicines)</li> </ul>	
Folic acid fortification and/or supplementation to prevent Neural Tube Defects <sup>10,11</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Folic acid fortification of staple food e.g. flour</li> <li>Folic acid tablets</li> </ul>	<ul style="list-style-type: none"> <li>Folic Acid for the Prevention of Neural Tube Defects: U.S. Preventive Services Task Force Recommendation Statement <a href="http://www.annals.org/content/150/9/626.abstract">www.annals.org/content/150/9/626.abstract</a></li> </ul>
<b>PREGNANCY</b>				
Antenatal Care <sup>12</sup> Essential Package	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Fetal stethoscope</li> <li>Scale</li> <li>Sphygmomanometer</li> <li>Haemoglobinometer</li> </ul>	<ul style="list-style-type: none"> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> <li>WHO Antenatal Care Randomized Trial: Manual for the implementation of the new model <a href="http://whqlibdoc.who.int/hq/2001/WHO_RHR_01.30.pdf">http://whqlibdoc.who.int/hq/2001/WHO_RHR_01.30.pdf</a></li> </ul>
Iron and folic acid supplementation during pregnancy <sup>13-15</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Iron and folic acid</li> </ul>	<ul style="list-style-type: none"> <li>Guidelines for the use of iron supplements to prevent and treat iron deficiency anaemia <a href="http://www.who.int/nutrition/publications/micronutrients/guidelines_for_Iron_supplementation.pdf">www.who.int/nutrition/publications/micronutrients/guidelines_for_Iron_supplementation.pdf</a></li> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> </ul>
Tetanus immunization in pregnancy for preventing neonatal tetanus <sup>16,17</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Vaccine (TT vaccine)</li> </ul>	<ul style="list-style-type: none"> <li>Neonatal tetanus <a href="http://www.who.int/immunization_monitoring/diseases/neonatal_tetanus/en/index.html">www.who.int/immunization_monitoring/diseases/neonatal_tetanus/en/index.html</a></li> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> </ul>

PRIORITY INTERVENTIONS	LEVEL OF CARE (COMMUNITY, PRIMARY, REFERRAL)	COMMUNITY OR PROFESSIONAL HEALTH WORKERS	KEY COMMODITIES (SUPPLEMENTED BY ANNEX)	PRACTICE GUIDELINES AND TRAINING MANUALS
<b>Prevention and management of malaria in pregnancy</b> a) Prophylactic antimalarial for preventing malaria in pregnancy <sup>18, 19</sup> b) Provision and promotion of use of Insecticide Treated Nets for preventing malaria in pregnancy <sup>20</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Antimalarial drugs according to the situation/context</li> <li>Insecticide Treated Nets</li> </ul>	<ul style="list-style-type: none"> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> <li>Insecticide treated bednets: a WHO position statement <a href="http://www.who.int/malaria/publications/atoz/itnspopaperfinal.pdf">www.who.int/malaria/publications/atoz/itnspopaperfinal.pdf</a></li> </ul>
Interventions for smoking cessation during pregnancy for improving birth outcomes <sup>21</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Materials for individual and group counselling and behavioural change interventions on smoking cessation</li> </ul>	<ul style="list-style-type: none"> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> </ul>
Screening and treatment of Syphilis <sup>22, 23</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Onsite tests and laboratory equipment</li> <li>Penicillin</li> <li>Counselling material</li> </ul>	<ul style="list-style-type: none"> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> <li>The Prevention and management of congenital syphilis: an overview and recommendations <a href="http://www.who.int/bulletin/volumes/82/6/424.pdf">www.who.int/bulletin/volumes/82/6/424.pdf</a></li> </ul>
<b>Prevention and management of HIV and Prevention of Mother-to-Child Transmission in Pregnancy</b> <sup>8, 24, 25</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>HIV test kits</li> <li>Antiretroviral drugs</li> <li>Cotrimoxazole</li> <li>Counselling material</li> </ul>	<ul style="list-style-type: none"> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> <li>Rapid advice: use of antiretroviral drugs for treating pregnant women and preventing HIV infection in infants <a href="http://www.who.int/hiv/pub/mtct/advice/en/index.html">www.who.int/hiv/pub/mtct/advice/en/index.html</a></li> </ul>
<b>Prevention and management of hypertension in pregnancy:</b> a) Calcium supplementation in pregnancy <sup>26-28</sup>	a) Community Primary Referral	a) ALL	a) Calcium	<ul style="list-style-type: none"> <li>WHO recommendations for the prevention and treatment of pre-eclampsia and eclampsia <a href="http://whqlibdoc.who.int/publications/2011/9789241548335_eng.pdf">http://whqlibdoc.who.int/publications/2011/9789241548335_eng.pdf</a></li> <li>Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> </ul>
b) Low-dose Aspirin for the prevention of pre-eclampsia in high risk women <sup>28, 29</sup>	b) Primary Referral	b) Professional health workers	b) Low dose Aspirin	
c) Use of antihypertensive drugs for treating severe hypertension in pregnancy <sup>28, 30</sup>	c) Primary Referral	c) Professional health workers	c) Methyldopa, Hydralazine, Nifedipine	
d) Prevention and treatment of Eclampsia <sup>28, 31, 32</sup>	d) Primary Referral	d) Professional health workers	d) Magnesium Sulphate (Injection)	
Reduce malpresentation at term using External Cephalic Version (> 36 weeks) <sup>33-35</sup>	Referral	Professional health workers	<ul style="list-style-type: none"> <li>Stethoscope</li> </ul>	<ul style="list-style-type: none"> <li>Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> </ul>



PRIORITY INTERVENTIONS	LEVEL OF CARE (COMMUNITY, PRIMARY, REFERRAL)	COMMUNITY OR PROFESSIONAL HEALTH WORKERS	KEY COMMODITIES (SUPPLEMENTED BY ANNEX)	PRACTICE GUIDELINES AND TRAINING MANUALS
<b>Management of prelabour rupture of membranes and preterm labour:</b>				
a) Induction of labour for management of prelabour rupture of membranes at term <sup>36</sup>	Referral	Professional health workers	<ul style="list-style-type: none"> <li>• Uterotonic (Oxytocin and/or Misoprostol)</li> <li>• Partograph</li> <li>• Stethoscope</li> <li>• Sphygmomanometer</li> </ul>	<ul style="list-style-type: none"> <li>• Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> <li>• WHO recommendations for induction of labour <a href="http://whqlibdoc.who.int/publications/2011/9789241501156_eng.pdf">http://whqlibdoc.who.int/publications/2011/9789241501156_eng.pdf</a></li> </ul>
b) Antibiotics for management of preterm rupture of membranes <sup>37</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>• Antibiotic (Erythromycin)</li> </ul>	<ul style="list-style-type: none"> <li>• Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> <li>• Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> </ul>
c) Corticosteroids for prevention of neonatal respiratory distress syndrome <sup>38, 39</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>• Corticosteroids (Betamethasone, Dexamethasone)</li> </ul>	<ul style="list-style-type: none"> <li>• Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> </ul>
<b>Management of unintended pregnancy:<sup>40</sup></b>	Primary Referral	Professional health workers		
a) Availability and provision of safe abortion			<ul style="list-style-type: none"> <li>• Materials for counselling, health education and health promotion</li> <li>• Medications for induced abortion (Mifepristone, Misoprostol)</li> <li>• Vacuum aspiration equipment</li> <li>• Uterotonics (Misoprostol, Oxytocin)</li> <li>• Antibiotics in line with essential medicine guidelines</li> <li>• Surgical procedures when required</li> <li>• Sphygmomanometer</li> </ul>	<ul style="list-style-type: none"> <li>• Safe abortion: technical and policy guidance for health systems. Geneva, World Health Organization, 2003 <a href="http://whqlibdoc.who.int/publications/2003/9241590343.pdf">http://whqlibdoc.who.int/publications/2003/9241590343.pdf</a></li> <li>• World Health Organization: Clinical practice handbook for safe abortion care. World Health Organization. Geneva. 2011. In Press</li> <li>• Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> </ul>
b) Provision of post abortion care				
<b>CHILDBIRTH</b>				
Social support during childbirth <sup>41</sup>	Community Primary Referral	ALL		<ul style="list-style-type: none"> <li>• Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> </ul>
Prophylactic antibiotic for caesarean section <sup>42</sup>	Referral	Professional health workers	<ul style="list-style-type: none"> <li>• Antibiotics (Ampicillin or Cefazolin)</li> </ul>	<ul style="list-style-type: none"> <li>• Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> </ul>
Caesarean section for maternal/foetal indication (e.g. obstructed labour and central placenta previa) (established practice)	Referral	Professional health workers	<ul style="list-style-type: none"> <li>• Surgical environment</li> <li>• Sphygmomanometer</li> </ul>	<ul style="list-style-type: none"> <li>• Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> </ul>
<b>Prevention of postpartum haemorrhage</b>				
a) Prophylactic uterotonic to prevent postpartum haemorrhage <sup>43, 44</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>• Uterotonics (Oxytocin, Misoprostol)</li> </ul>	<ul style="list-style-type: none"> <li>• WHO recommendation for prevention of postpartum haemorrhage <a href="http://whqlibdoc.who.int/hq/2007/WHO_MPS_07.06_eng.pdf">http://whqlibdoc.who.int/hq/2007/WHO_MPS_07.06_eng.pdf</a></li> </ul>



PRIORITY INTERVENTIONS	LEVEL OF CARE (COMMUNITY, PRIMARY, REFERRAL)	COMMUNITY OR PROFESSIONAL HEALTH WORKERS	KEY COMMODITIES (SUPPLEMENTED BY ANNEX)	PRACTICE GUIDELINES AND TRAINING MANUALS
b) Active management of third stage of labour to prevent postpartum haemorrhage <sup>45-49</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Uterotonics (Oxytocin, Ergometrine)</li> </ul>	<ul style="list-style-type: none"> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> <li>Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> <li>WHO recommendation for prevention of postpartum haemorrhage <a href="http://whqlibdoc.who.int/hq/2007/WHO_MPS_07.06_eng.pdf">http://whqlibdoc.who.int/hq/2007/WHO_MPS_07.06_eng.pdf</a></li> </ul>
Induction of labour for prolonged pregnancy <sup>50</sup>	Referral	Professional health workers	<ul style="list-style-type: none"> <li>Uterotonics (Oxytocin, Misoprostol)</li> </ul>	<ul style="list-style-type: none"> <li>Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> <li>Managing prolonged and obstructed labour <a href="http://www.who.int/making_pregnancy_safer/documents/3_9241546662/en/index.html">www.who.int/making_pregnancy_safer/documents/3_9241546662/en/index.html</a></li> <li>WHO recommendations for induction of labour <a href="http://whqlibdoc.who.int/publications/2011/9789241501156_eng.pdf">http://whqlibdoc.who.int/publications/2011/9789241501156_eng.pdf</a></li> </ul>
<b>Management of postpartum haemorrhage</b> e.g. a) uterine massage b) uterotonics <sup>48, 51</sup>	Community Primary Referral	Community health workers Primary and Referral	<ul style="list-style-type: none"> <li>Uterotonics (Oxytocin, Ergometrine, Misoprostol)</li> </ul>	<ul style="list-style-type: none"> <li>Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> <li>WHO guidelines for the management of postpartum haemorrhage and retained placenta <a href="http://whqlibdoc.who.int/publications/2009/9789241598514_eng.pdf">http://whqlibdoc.who.int/publications/2009/9789241598514_eng.pdf</a></li> </ul>
c) manual removal of placenta (only by professional health workers)	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Uterotonics (Oxytocin, Ergometrine, Misoprostol)</li> <li>IV fluids</li> <li>Blood transfusion</li> <li>Surgical facilities</li> </ul>	
Initiation or continuation of HIV therapy for HIV positive women	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>HIV testing kit + ARVs</li> </ul>	<ul style="list-style-type: none"> <li>Rapid Advice: Use of antiretroviral drugs for treating pregnant women and preventing HIV infection in infants <a href="http://whqlibdoc.who.int/publications/2009/9789241598934_eng.pdf">http://whqlibdoc.who.int/publications/2009/9789241598934_eng.pdf</a></li> </ul>
<b>POSTNATAL - MOTHER</b>				
Advice and provision of family planning <sup>52</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Barrier methods (male and female condoms, diaphragm, gels, foams)</li> <li>Oral contraceptives (progestin only and combined)</li> <li>Emergency contraception and hormonal injections</li> </ul>	<ul style="list-style-type: none"> <li>Medical eligibility criteria for contraceptive use <a href="http://whqlibdoc.who.int/publications/2010/9789241563888_eng.pdf">http://whqlibdoc.who.int/publications/2010/9789241563888_eng.pdf</a></li> <li>Family Planning: a global handbook for providers <a href="http://whqlibdoc.who.int/publications/2011/9780978856373_eng.pdf">http://whqlibdoc.who.int/publications/2011/9780978856373_eng.pdf</a></li> </ul>
	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>All of the above plus implants</li> <li>Long acting reversible contraceptives (implants)</li> <li>Intrauterine devices</li> <li>Surgical contraception</li> </ul>	<ul style="list-style-type: none"> <li>Surgical Care at the District Hospital <a href="http://www.who.int/surgery/publications/scdh_manual/en/index.html">www.who.int/surgery/publications/scdh_manual/en/index.html</a> pgs 9-8, 11-19</li> </ul>
Prevent, measure and treat maternal anaemia <sup>53</sup>	Referral	Professional health workers	<ul style="list-style-type: none"> <li>Ferrous Salt (liquid or tablet)</li> <li>Ferrous Salt+Folic Acid (tablet)</li> <li>Folic Acid (tablet)</li> <li>Hydroxycobalamine (injection)</li> <li>Lab tests</li> <li>Blood products</li> </ul>	<ul style="list-style-type: none"> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> <li>Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> </ul>

PRIORITY INTERVENTIONS	LEVEL OF CARE (COMMUNITY, PRIMARY, REFERRAL)	COMMUNITY OR PROFESSIONAL HEALTH WORKERS	KEY COMMODITIES (SUPPLEMENTED BY ANNEX)	PRACTICE GUIDELINES AND TRAINING MANUALS
Detection and management of postpartum sepsis <sup>54</sup>	Referral	Professional health workers	<ul style="list-style-type: none"> <li>Antibiotics (Ampicillin, Gentamicin, Metronidazole)</li> </ul>	<ul style="list-style-type: none"> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> <li>Managing Complications in Pregnancy and Childbirth: A guide for midwives and doctors <a href="http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf">http://whqlibdoc.who.int/publications/2007/9241545879_eng.pdf</a></li> </ul>
Screening and initiation or continuation of antiretroviral therapy for HIV <sup>8</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Antiretroviral medicines</li> <li>HIV test kits</li> </ul>	<ul style="list-style-type: none"> <li>Pregnancy, Childbirth, Postpartum and Newborn Care: a guide to essential practice <a href="http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf">http://whqlibdoc.who.int/publications/2006/924159084X_eng.pdf</a></li> <li>Rapid advice: use of antiretroviral drugs for treating pregnant women and preventing HIV infection in infants <a href="http://www.who.int/hiv/pub/mtct/advice/en/index.html">www.who.int/hiv/pub/mtct/advice/en/index.html</a></li> </ul>



# NEWBORN CARE INTERVENTIONS - BIRTH AND POSTNATAL

PRIORITY INTERVENTIONS	LEVEL OF CARE (COMMUNITY, PRIMARY, REFERRAL)	COMMUNITY OR PROFESSIONAL HEALTH WORKERS	KEY COMMODITIES (SUPPLEMENTED BY ANNEX)	PRACTICE GUIDELINES AND TRAINING MANUALS
<b>IMMEDIATE ESSENTIAL NEWBORN CARE (AT THE TIME OF BIRTH)</b>				
Promotion and provision of thermal care for all newborns to prevent hypothermia (immediate drying, warming, skin to skin, delayed bathing) <sup>55</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Materials for counselling, health education and health promotion</li> </ul>	<ul style="list-style-type: none"> <li>WHO essential newborn care <a href="http://www.who.int/making_pregnancy_safer/documents/newborncare_course/en/index.html">www.who.int/making_pregnancy_safer/documents/newborncare_course/en/index.html</a></li> <li>WHO. Thermal protection of the newborn: a practical guide (Part of training material) <a href="http://www.who.int/making_pregnancy_safer/documents/ws42097th/en/">http://www.who.int/making_pregnancy_safer/documents/ws42097th/en/</a></li> <li>WHO. IMCI chart booklet (2008) <a href="http://www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html">www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html</a></li> </ul>
Promotion and support for early initiation and exclusive breastfeeding (within the first hour) <sup>56-59</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Materials for counselling, health education and health promotion</li> </ul>	<ul style="list-style-type: none"> <li>WHO. Infant and Young child feeding - Programming Guide <a href="http://www.who.int/child_adolescent_health/documents/9241591218/en/index.html">www.who.int/child_adolescent_health/documents/9241591218/en/index.html</a></li> <li>WHO. IMCI chart booklet (2008) <a href="http://www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html">www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html</a></li> <li>Infant young child feeding counselling: An integrated course (Part of training material) <a href="http://www.who.int/nutrition/publications/infantfeeding/9789241594745/en/index.html">www.who.int/nutrition/publications/infantfeeding/9789241594745/en/index.html</a></li> </ul>
Promotion and provision of hygienic cord and skin care <sup>60</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Cord clamp and scissors</li> <li>Clean birth kit for health facilities</li> </ul>	<ul style="list-style-type: none"> <li>WHO Essential newborn care <a href="http://www.who.int/making_pregnancy_safer/documents/newborncare_course/en/index.html">www.who.int/making_pregnancy_safer/documents/newborncare_course/en/index.html</a></li> <li>WHO. IMCI chart booklet (2008) <a href="http://www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html">www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html</a></li> <li>WHO. IMPAC - Pregnancy, childbirth, postpartum and newborn care: a guide for essential practice (2006) <a href="http://www.who.int/making_pregnancy_safer/documents/924159084x/en/index.html">www.who.int/making_pregnancy_safer/documents/924159084x/en/index.html</a></li> </ul>
Neonatal resuscitation with bag and mask for babies who do not breathe at birth <sup>61-63</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Training aids and devices to maintain competencies</li> <li>Newborn resuscitation device (Ambu Bag, bag-mask and suction device)</li> </ul>	<ul style="list-style-type: none"> <li>American Academy of Pediatrics Helping babies breathe - The Golden Minute <a href="http://www.helpingbabiesbreathe.org/masterTrainers.html">www.helpingbabiesbreathe.org/masterTrainers.html</a></li> <li>WHO Essential newborn care <a href="http://www.who.int/making_pregnancy_safer/documents/newborncare_course/en/index.html">www.who.int/making_pregnancy_safer/documents/newborncare_course/en/index.html</a></li> </ul>
Newborn immunization	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Vaccines, syringes, safety boxes, cold chain equipment</li> </ul>	<ul style="list-style-type: none"> <li>WHO Vaccine Position papers <a href="http://www.who.int/immunization/position_papers/en/">www.who.int/immunization/position_papers/en/</a></li> </ul>
<b>NEONATAL INFECTION MANAGEMENT</b>				
Presumptive antibiotic therapy for the newborns at risk of bacterial infection <sup>64</sup>	Referral	Professional health workers	<ul style="list-style-type: none"> <li>Antibiotics (ampicillin and gentamicin or penicillin)</li> </ul>	<ul style="list-style-type: none"> <li>WHO. Managing newborn problems - a guide for doctors, nurses and midwives <a href="http://www.who.int/making_pregnancy_safer/documents/9241546220/en/index.html">www.who.int/making_pregnancy_safer/documents/9241546220/en/index.html</a></li> <li>WHO. IMPAC - Pregnancy, childbirth, postpartum and newborn care: a guide for essential practice (2006) <a href="http://www.who.int/making_pregnancy_safer/documents/924159084x/en/index.html">www.who.int/making_pregnancy_safer/documents/924159084x/en/index.html</a></li> </ul>
Case management of neonatal sepsis, meningitis and pneumonia <sup>65-69</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Materials for counselling, health education and health promotion</li> <li>Thermometer / digital thermometer</li> <li>Timer</li> <li>Blood sugar sticks (disposable)</li> <li>Nasogastric tube</li> <li>Antibiotics (oral and injectable)</li> </ul>	<ul style="list-style-type: none"> <li>WHO. IMCI chart booklet (2008) <a href="http://www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html">www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html</a></li> <li>WHO. Pocket book of hospital care for children (2005) <a href="http://www.who.int/child_adolescent_health/documents/9241546700/en/index.html">www.who.int/child_adolescent_health/documents/9241546700/en/index.html</a></li> </ul>
Initiation of ART in babies born to HIV infected mother	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>HIV testing kit + ARVs</li> </ul>	<ul style="list-style-type: none"> <li>Rapid Advice: Use of antiretroviral drugs for treating pregnant women and preventing HIV infection in infants <a href="http://whqlibdoc.who.int/publications/2009/9789241598934_eng.pdf">http://whqlibdoc.who.int/publications/2009/9789241598934_eng.pdf</a></li> </ul>



PRIORITY INTERVENTIONS	LEVEL OF CARE (COMMUNITY, PRIMARY, REFERRAL)	COMMUNITY OR PROFESSIONAL HEALTH WORKERS	KEY COMMODITIES (SUPPLEMENTED BY ANNEX)	PRACTICE GUIDELINES AND TRAINING MANUALS
<b>INTERVENTIONS FOR SMALL AND ILL BABIES</b>				
Kangaroo mother care (KMC) for preterm and for < 2000g babies <sup>70, 71</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Materials for counselling, health education and health promotion</li> <li>Support Binder for KMC (KMC wrap)</li> <li>Hat</li> <li>Nasogastric tube</li> </ul>	<ul style="list-style-type: none"> <li>WHO   Kangaroo mother care: a practical guide <a href="http://www.who.int/making_pregnancy_safer/documents/9241590351/en/">www.who.int/making_pregnancy_safer/documents/9241590351/en/</a></li> <li>WHO. Essential newborn care course (2010) - Training Tool <a href="http://www.who.int/making_pregnancy_safer/documents/newborn_care_course/en/">www.who.int/making_pregnancy_safer/documents/newborn_care_course/en/</a></li> </ul>
Extra support for feeding the small and preterm baby <sup>72</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Nasogastric tubes</li> <li>Feeding cups</li> <li>Breast pump</li> <li>Syringe drivers</li> <li>Blood sugar testing sticks</li> <li>Materials for counselling</li> </ul>	<ul style="list-style-type: none"> <li>WHO guide for feeding preterm and LBW babies (forthcoming in the web)</li> <li>WHO. Essential newborn care course (2010) - Training Tool <a href="http://www.who.int/making_pregnancy_safer/documents/newborn_care_course/en/">www.who.int/making_pregnancy_safer/documents/newborn_care_course/en/</a></li> </ul>
Prophylactic and therapeutic use of surfactant to prevent respiratory distress syndrome in pre-term babies <sup>73</sup>	Referral	Professional health workers	<ul style="list-style-type: none"> <li>Surfactant</li> <li>Oxygen supply/concentrator</li> <li>Pulse oximeter</li> </ul>	<ul style="list-style-type: none"> <li>WHO. IMPAC - Managing newborn problems: a guide for doctors, nurses and midwives (2003) - Guideline <a href="http://www.who.int/making_pregnancy_safer/documents/9241546220/en/index.html">www.who.int/making_pregnancy_safer/documents/9241546220/en/index.html</a></li> </ul>
Continuous positive airway pressure (CPAP) to manage pre-term babies with respiratory distress syndrome <sup>74, 75</sup>	Referral	Professional health workers	<ul style="list-style-type: none"> <li>Standard CPAP or bubble CPAP</li> <li>Oxygen supply/concentrator</li> <li>Pulse oximeter</li> </ul>	<ul style="list-style-type: none"> <li>WHO. IMPAC - Managing newborn problems: a guide for doctors, nurses and midwives (2003) <a href="http://www.who.int/making_pregnancy_safer/documents/9241546220/en/index.html">www.who.int/making_pregnancy_safer/documents/9241546220/en/index.html</a></li> </ul>
Management of newborns with jaundice <sup>76, 77</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Bilirubinometer</li> <li>Phototherapy lamp</li> <li>eye shade</li> <li>IV fluids</li> <li>Exchange transfusion kit</li> </ul>	<ul style="list-style-type: none"> <li>WHO. Pocket book of hospital care for children (2005) <a href="http://www.who.int/child_adolescent_health/documents/9241546700/en/index.html">www.who.int/child_adolescent_health/documents/9241546700/en/index.html</a></li> <li>WHO. IMPAC - Managing newborn problems: a guide for doctors, nurses and midwives (2003) <a href="http://www.who.int/making_pregnancy_safer/documents/9241546220/en/index.html">www.who.int/making_pregnancy_safer/documents/9241546220/en/index.html</a></li> </ul>



PRIORITY INTERVENTIONS	LEVEL OF CARE (COMMUNITY, PRIMARY, REFERRAL)	COMMUNITY OR PROFESSIONAL HEALTH WORKERS	KEY COMMODITIES (SUPPLEMENTED BY ANNEX)	PRACTICE GUIDELINES AND TRAINING MANUALS
<b>INFANCY AND CHILDHOOD</b>				
Promotion and support for exclusive breastfeeding for 6 months <sup>78, 79</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Materials for counselling, health education and health promotion, including individual and group counselling</li> </ul>	<ul style="list-style-type: none"> <li>WHO. Exclusive Breastfeeding <a href="http://www.who.int/nutrition/topics/exclusive_breastfeeding/en/">www.who.int/nutrition/topics/exclusive_breastfeeding/en/</a></li> <li>WHO. Infant and young child feeding counselling: an integrated course (2006) - Training tool <a href="http://www.who.int/nutrition/publications/infantfeeding/9789241594745/en/index.html">www.who.int/nutrition/publications/infantfeeding/9789241594745/en/index.html</a></li> <li>WHO. Community-based strategies for breastfeeding promotion and support in developing countries (2003) - Technical Review <a href="http://www.who.int/child_adolescent_health/documents/9241591218/en/index.html">www.who.int/child_adolescent_health/documents/9241591218/en/index.html</a></li> <li>WHO. IMCI chart booklet (2008) <a href="http://www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html">www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html</a></li> </ul>
<b>Promotion and support of continued breastfeeding and complementary feeding</b> <ol style="list-style-type: none"> <li>Continued breastfeeding up to 2 years and beyond<sup>79</sup></li> <li>Appropriate complementary feeding starting at 6 months<sup>80, 81</sup></li> </ol>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Materials for counselling, health education and health promotion</li> </ul>	<ul style="list-style-type: none"> <li>WHO. Guiding principles for complementary feeding of the breastfed child (2003) <a href="http://whqlibdoc.who.int/paho/2003/a85622.pdf">http://whqlibdoc.who.int/paho/2003/a85622.pdf</a></li> <li>WHO. Guiding principles for feeding non-breastfed children 6-24 months of age (2005) <a href="http://www.who.int/child_adolescent_health/documents/9241593431/en/index.html">www.who.int/child_adolescent_health/documents/9241593431/en/index.html</a></li> </ul>
<b>Prevention and management of childhood malaria</b> <ol style="list-style-type: none"> <li>Provision and promotion of use of insecticide treated bed nets for children<sup>82, 83</sup></li> <li>Case management of childhood malaria<sup>84</sup></li> </ol>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Materials for counselling, health education and health promotion</li> <li>Insecticide treated nets</li> <li>Rapid diagnostic tests</li> <li>Antimalarial drugs according to guidelines</li> </ul>	<ul style="list-style-type: none"> <li>WHO. Insecticide-treated mosquito nets: a position statement (2007) <a href="http://www.who.int/malaria/publications/atoz/itnspospaperfinal/en/index.html">www.who.int/malaria/publications/atoz/itnspospaperfinal/en/index.html</a></li> <li>WHO. Guidelines for the treatment of malaria (2010) <a href="http://whqlibdoc.who.int/publications/2010/9789241547925_eng.pdf">http://whqlibdoc.who.int/publications/2010/9789241547925_eng.pdf</a></li> <li>WHO. IMCI chart booklet being updated at <a href="http://www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html">www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html</a></li> <li>WHO. Pocket book of hospital care for children: guidelines for the management of common illnesses with limited resources (being updated) <a href="http://www.who.int/child_adolescent_health/documents/9241546700/en/index.html">www.who.int/child_adolescent_health/documents/9241546700/en/index.html</a></li> <li>WHO. Emergency Triage Assessment and Treatment (ETAT) course at <a href="http://www.who.int/child_adolescent_health/documents/9241546875/en/index.html">www.who.int/child_adolescent_health/documents/9241546875/en/index.html</a></li> </ul>

PRIORITY INTERVENTIONS	LEVEL OF CARE (COMMUNITY, PRIMARY, REFERRAL)	COMMUNITY OR PROFESSIONAL HEALTH WORKERS	KEY COMMODITIES (SUPPLEMENTED BY ANNEX)	PRACTICE GUIDELINES AND TRAINING MANUALS
Comprehensive care of children infected with or exposed to HIV <sup>8, 85</sup>	Primary Referral	Professional health workers	<ul style="list-style-type: none"> <li>Antiretroviral drugs</li> <li>HIV test kits</li> <li>Cotrimoxazole</li> <li>Psychosocial support</li> <li>Nutritional support</li> </ul>	<ul style="list-style-type: none"> <li>WHO. Guidelines on HIV and infant feeding 2010 <a href="http://www.who.int/nutrition/publications/hivaids/9789241599535/en/index.html">www.who.int/nutrition/publications/hivaids/9789241599535/en/index.html</a></li> <li>WHO. Manual on paediatric HIV care and treatment for district hospitals <a href="http://www.who.int/child_adolescent_health/documents/9789241501026/en/index.html">www.who.int/child_adolescent_health/documents/9789241501026/en/index.html</a></li> <li>WHO recommendations on the management of diarrhoea and pneumonia in HIV-infected infants and children <a href="http://www.who.int/child_adolescent_health/documents/9789241548083/en/index.html">www.who.int/child_adolescent_health/documents/9789241548083/en/index.html</a></li> <li>WHO. IMCI chart booklet for high HIV settings <a href="http://www.who.int/child_adolescent_health/documents/9789241597388/en/index.html">www.who.int/child_adolescent_health/documents/9789241597388/en/index.html</a></li> <li>WHO. Pocket book of hospital care for children: guidelines for the management of common illnesses with limited resources <a href="http://www.who.int/child_adolescent_health/documents/9241546700/en/index.html">www.who.int/child_adolescent_health/documents/9241546700/en/index.html</a></li> </ul>
Promote and provide routine immunization plus <i>H.influenzae</i> , meningococcal, pneumococcal, and rotavirus vaccines <sup>86, 87</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Materials for counselling, health education and health promotion</li> <li>Vaccines, syringes, safety boxes, cold chain equipment</li> </ul>	<ul style="list-style-type: none"> <li>WHO. IMCI chart booklet (2008) - Guideline <a href="http://www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html">www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html</a></li> </ul>
Vitamin A supplementation from 6 months of age in Vitamin A deficient populations <sup>88, 89</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>Vitamin A capsules</li> <li>Material for counselling on Vitamin A rich foods</li> </ul>	<ul style="list-style-type: none"> <li>WHO Guideline: Vitamin A supplementation in infants and children 6-59 months of age (2011) <a href="http://www.who.int/nutrition/publications/micronutrients/guidelines/vas_6to59_months/en/index.html">www.who.int/nutrition/publications/micronutrients/guidelines/vas_6to59_months/en/index.html</a></li> </ul>
<b>Management of severe acute malnutrition:</b> <sup>90, 91</sup> a) without complications (all levels) b) with complications (Referral)	Community Primary Referral	ALL	<p><b>Community level</b></p> <ul style="list-style-type: none"> <li>Appropriate ready-to-use therapeutic foods</li> <li>Micronutrient supplements</li> <li>Vitamin A capsules</li> </ul> <p><b>Health Facility level</b></p> <ul style="list-style-type: none"> <li>Antibiotics</li> <li>Therapeutic food formulations (F75/100)</li> </ul>	<ul style="list-style-type: none"> <li>WHO. Management of severe malnutrition: a manual for physicians and other senior health workers (1999) <a href="http://www.who.int/nutrition/publications/severemalnutrition/en/manage_severe_malnutrition_eng.pdf">www.who.int/nutrition/publications/severemalnutrition/en/manage_severe_malnutrition_eng.pdf</a></li> <li>WHO. Pocket book of hospital care for children: guidelines for the management of common illnesses with limited resources <a href="http://www.who.int/child_adolescent_health/documents/9241546700/en/index.html">www.who.int/child_adolescent_health/documents/9241546700/en/index.html</a></li> </ul>
<b>Case management of childhood pneumonia</b> <sup>92</sup> a) Vitamin A as part of treatment for measles-associated pneumonia for children above 6 months <sup>93, 94</sup> b) Vitamin A as part of treatment for non-measles-associated pneumonia for children above 6 months <sup>92, 95-98</sup>	Community Primary Referral	ALL	<p><b>Community and Health Facility level</b></p> <ul style="list-style-type: none"> <li>Respiratory rate timers</li> <li>Vitamin A capsules</li> <li>Appropriate antibiotics</li> </ul> <p><b>Referral level</b></p> <ul style="list-style-type: none"> <li>Oxygen for severe pneumonia</li> <li>Pulse oximeter</li> </ul>	<ul style="list-style-type: none"> <li>WHO. Manual for the Community Health Worker: Caring for the sick child in the community (Working Version)</li> <li>WHO and UNICEF. Management of Sick Children by Community Health Worker (2006) <a href="http://www.unicef.org/publications/files/Management_of_Sick_Children_by_Community_Health_Workers.pdf">www.unicef.org/publications/files/Management_of_Sick_Children_by_Community_Health_Workers.pdf</a></li> <li>WHO. IMCI chart booklet (2008) - Guideline <a href="http://www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html">www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html</a></li> <li>WHO. Pocket book of hospital care for children - Guideline <a href="http://www.who.int/child_adolescent_health/documents/9241546700/en/index.html">www.who.int/child_adolescent_health/documents/9241546700/en/index.html</a></li> </ul>



PRIORITY INTERVENTIONS	LEVEL OF CARE (COMMUNITY, PRIMARY, REFERRAL)	COMMUNITY OR PROFESSIONAL HEALTH WORKERS	KEY COMMODITIES (SUPPLEMENTED BY ANNEX)	PRACTICE GUIDELINES AND TRAINING MANUALS
<b>Case management of diarrhoea:</b> a) Acute watery diarrhoea <sup>99-105</sup> b) Dysentery <sup>106, 107</sup>	Community Primary Referral	ALL	<ul style="list-style-type: none"> <li>• Materials for counselling, health education and health promotion</li> <li>• Zinc (tablets / solution)</li> <li>• Oral Rehydration Solution (ORS)</li> <li>• Appropriate antibiotics for dysentery according to guidelines</li> </ul>	<ul style="list-style-type: none"> <li>• WHO Guidelines on hand hygiene in health care (2009) <a href="http://whqlibdoc.who.int/publications/2009/9789241597906_eng.pdf">http://whqlibdoc.who.int/publications/2009/9789241597906_eng.pdf</a></li> <li>• WHO. Guidelines for Drinking Water Safety (2011) <a href="http://www.who.int/water_sanitation_health/publications/2011/dwq_guidelines/en/index.html">www.who.int/water_sanitation_health/publications/2011/dwq_guidelines/en/index.html</a></li> <li>• WHO. Guidelines for the safe use of wastewater, excreta and greywater (2006) <a href="http://www.who.int/water_sanitation_health/wastewater/gsuww/en/index.html">www.who.int/water_sanitation_health/wastewater/gsuww/en/index.html</a></li> <li>• WHO. Manual for the Community Health Worker: Caring for the sick child in the community (Working Version)</li> <li>• WHO. Management of Sick Children by Community Health Worker (2006) <a href="http://www.unicef.org/publications/files/Management_of_Sick_Children_by_Community_Health_Workers.pdf">www.unicef.org/publications/files/Management_of_Sick_Children_by_Community_Health_Workers.pdf</a></li> <li>• WHO. IMCI chart booklet (2008) - Guideline <a href="http://www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html">www.who.int/child_adolescent_health/documents/IMCI_chartbooklet/en/index.html</a></li> <li>• WHO. Pocket book of hospital care for children - Guideline <a href="http://www.who.int/child_adolescent_health/documents/9241546700/en/index.html">www.who.int/child_adolescent_health/documents/9241546700/en/index.html</a></li> </ul>
Case management of meningitis	Referral	Professional health workers	<ul style="list-style-type: none"> <li>• Appropriate antibiotics</li> <li>• Supportive treatment</li> </ul>	<ul style="list-style-type: none"> <li>• WHO. Pocket book of hospital care for children - Guideline <a href="http://www.who.int/child_adolescent_health/documents/9241546700/en/index.html">www.who.int/child_adolescent_health/documents/9241546700/en/index.html</a></li> </ul>



## CROSS-CUTTING COMMUNITY STRATEGIES

There are several cross-cutting community strategies across the continuum of care e.g. home visits<sup>59, 108</sup>, participation of women's groups, conditional cash transfers etc. The strategies can be used for demand creation, empowerment, service delivery etc. These will be reviewed at a later date and recommendations arising from the review will be published in a complementary document.





1. Trends in Maternal Mortality 1990 to 2008. Estimates developed by WHO, UNICEF, UNFPA and The World Bank. (2010)
2. Levels & trends in Child Mortality. Report 2011. Estimates Developed by the UN Inter-agency Group for Child Mortality Estimation. (2011)
3. Bhutta Z, Yakoob M, Salam R, Lassi Z. Global Review of Interventions Related to Maternal, Newborn and Child Health (MNCH): What Works and Can be Scaled-up? Aga Khan University. Pakistan. 2011. Available on [www.pmnch.org](http://www.pmnch.org)
4. Jamison D, Breman G, Measham A, Alleyne G, Claeson M, Evans, D; Jha, P; Mills, A; Musgrove, P. Priorities in Health. The World Bank. April 2006. Washington D.C.
5. Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Birth spacing and risk of adverse perinatal outcomes. *JAMA*. 2006;295(15):1809-1823.
6. Conde-Agudelo A, Belizan JM, Breman R, Brockman SC, Rosas-Bermudez A. Effect of the interpregnancy interval after an abortion on maternal and perinatal health in Latin America. *International Journal of Gynecology & Obstetrics*. 2005;89:S34-S40.
7. Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Effects of birth spacing on maternal health: a systematic review. *American Journal of Obstetrics and Gynecology*. 2007;196(4):297-308.
8. Siegfried N, van der Merwe L, Brocklehurst P, Sint TT. Antiretrovirals for reducing the risk of mother-to-child transmission of HIV infection. *Cochrane Database of Systematic Reviews*. 2011; Issue 7. Art. No.: CD003510.
9. Ng BE, Butler LM, Horvath T, Rutherford GW. Population-based biomedical sexually transmitted infection control interventions for reducing HIV infection. *Cochrane Database of Systematic Reviews*. 2011; Issue 3. Art. No.: CD001220.
10. De-Regil LM, Fernández-Gaxiola AC, Dowswell T, Peña-Rosas JP. Effects and safety of periconceptional folate supplementation for preventing birth defects. *Cochrane Database of Systematic Reviews*. 2010; Issue 10. Art. No.: CD007950.
11. Blencowe H, Cousens S, Modell B, Lawn J. Folic acid to reduce neonatal mortality from neural tube disorders. *International Journal of Epidemiology*. 2010;39(Suppl. 1):i110-i121.
12. Dowswell T, Carroli G, Duley L, et al. Alternative versus standard packages of antenatal care for low-risk pregnancy. *Cochrane Database of Systematic Reviews*. 2010; Issue 10. Art. No.: CD000934.
13. Peña-Rosas JP, Viteri FE. Effects and safety of preventive oral iron or iron+ folic acid supplementation for women during pregnancy. *Cochrane Database of Systematic Reviews*. 2009; Issue 4. Art. No.: CD005462.
14. Yakoob MY, Bhutta ZA. Effect of routine iron supplementation with or without folic acid on anaemia during pregnancy. *BMC Public Health*. 2011;11(Suppl 3):S21.
15. Imdad A, Yakoob MY, Bhutta ZA. The effect of folic acid, protein energy and multiple micronutrient supplements in pregnancy on stillbirths. *BMC Public Health*. 2011;11(Suppl 3):S4.
16. Demicheli V, Barale A, Rivetti A. Vaccines for women to prevent neonatal tetanus. *Cochrane Database of Systematic Reviews*. 2005; Issue 4. Art. No.: CD002959.
17. Blencowe H, Joy Lawn J, Vandelaer J, Roper M, Cousens S. Tetanus toxoid immunization to reduce mortality from neonatal tetanus. *International Journal of Epidemiology*. 2010;39:i102-i109.
18. Eisele TP, Larsen D, Steketee RW. Protective efficacy of interventions for preventing malaria mortality in children in Plasmodium falciparum endemic areas. *International Journal of Epidemiology*. 2010;39(suppl 1):i88-i101.
19. Garner P, Gulmezoglu AM. Drugs for preventing malaria in pregnant women. *Cochrane Database of Systematic Reviews*. 2006; Issue 4. Art. No.: CD000169.
20. Gamble C, Ekwaru JP, Ter Kuile FO. Insecticide-treated nets for preventing malaria in pregnancy. *Cochrane Database of Systematic Reviews*. 2006; Issue 2. Art. No.: CD003755.
21. Lumley J, Chamberlain C, Dowswell T, et al. Interventions for promoting smoking cessation during pregnancy. *Cochrane Database of Systematic Reviews*. 2009; Issue 3. Art. No.: CD001055.
22. Blencowe H, Cousens S, Kamb M, Berman S, Lawn JE. Lives Saved Tool supplement detection and treatment of syphilis in pregnancy to reduce syphilis related stillbirths and neonatal mortality. *BMC Public Health*. 2011;11(Suppl 3):S9:1-16.
23. Walker GJA. Antibiotics for syphilis diagnosed during pregnancy. *Cochrane Database of Systematic Reviews*. 2001; Issue 3. Art. No.: CD001143.
24. Read JS, Newell ML. Efficacy and safety of cesarean delivery for prevention of mother-to-child transmission of HIV-1. *Cochrane Database of Systematic Reviews*. 2005; Issue 4. Art. No.: CD005479.
25. Sturt AS, Dokubo EK, Sint TT. Antiretroviral therapy (ART) for treating HIV infection in ART-eligible pregnant women. *Cochrane Database of Systematic Reviews*. 2010; Issue 3. Art. No.: CD008440.
26. Hofmeyr GJ, Lawrie TA, Atallah AN, Duley L. Calcium supplementation during pregnancy for preventing hypertensive disorders and related problems. *Cochrane Database of Systematic Reviews*. 2010; Issue 8. Art. No.: CD001059.
27. Imdad A, Jabeen A, Bhutta Z. Role of calcium supplementation during pregnancy in reducing risk of developing gestational hypertensive disorders: a meta-analysis of studies from developing countries. *BMC Public Health*. 2011;11(Suppl 3):S18.
28. Jabeen M, Yakoob MY, Imdad A, Bhutta ZA. Impact of interventions to prevent and manage preeclampsia and eclampsia on stillbirths. *BMC Public Health*. 2011;11(Suppl 3):S6.
29. Duley L, Henderson-Smart DJ, Knight M, King JF. Antiplatelet agents for preventing pre-eclampsia and its complications. *Cochrane Database of Systematic Reviews*. 2007; Issue 2. Art. No.: CD004659.
30. Duley L, Henderson-Smart DJ, Meher S. Drugs for treatment of very high blood pressure during pregnancy. *Cochrane Database of Systematic Reviews*. 2006; Issue 3. Art. No.: CD001449.
31. Duley L, Henderson-Smart DJ, Chou D. Magnesium sulphate versus phenytoin for eclampsia. *Cochrane Database of Systematic Reviews*. 2010; Issue 12. Art. No.: CD000127.
32. Duley L, Henderson-Smart DJ, Walker GJ, Chou D. Magnesium sulphate versus diazepam for eclampsia. *Cochrane Database of Systematic Reviews*. 2010; Issue 10. Art. No.: CD000128.
33. Hofmeyr GJ, Gyte GML. Interventions to help external cephalic version for breech presentation at term. *Cochrane Database of Systematic Reviews*. 2004; Issue 1. Art. No.: CD000184.
34. Coyle ME, Smith CA, Peat B. Cephalic version by moxibustion for breech presentation. *Cochrane Database of Systematic Reviews*. 2005; Issue 2. Art. No.: CD003928.
35. Hofmeyr GJ, Kulier R. Cephalic version by postural management for breech presentation. *Cochrane Database of Systematic Reviews*. 2000; Issue 3. Art. No.: CD000051.
36. Kenyon S, Boulvain M, Neilson JP. Antibiotics for preterm rupture of membranes. *Cochrane Database of Systematic Reviews*. 2010; Issue 8. Art. No.: CD001058.
37. Cousens S, Blencowe H, Gravett M, Lawn JE. Antibiotics for pre-term pre-labour rupture of membranes: prevention of neonatal deaths due to complications of pre-term birth and infection. *International Journal of Epidemiology*. 2010;39(Suppl. 1):i134-i143.
38. Mwansa-Kambafwile J, Cousens S, Hansen T, Lawn JE. Antenatal steroids in preterm labour for the prevention of neonatal deaths due to complications of preterm birth. *International Journal of Epidemiology*. 2010;39(Suppl 1):i122-i133.

39. Roberts D, Dalziel SR. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database of Systematic Reviews*. 2006;Issue 3. Art. No.: CD004454.
40. World Health Organization. *Safe abortion: technical and policy guidance for health systems* Geneva World Health Organization; 2003.
41. Hodnett ED, Gates S, Hofmeyr GJ, Sakala C, Weston J. Continuous support for women during childbirth. *Cochrane Database of Systematic Reviews*. 2011;Issue 2. Art. No.: CD003766.
42. Smaill FM, Gyte GM. Antibiotic prophylaxis versus no prophylaxis for preventing infection after cesarean section. *Cochrane Database for Systematic Reviews*. 2010;Issue 1. Art. No.: CD007482.
43. Soltani H, Hutchon DR, Poulouse TA. Timing of prophylactic uterotonics for the third stage of labour after vaginal birth. *Cochrane Database of Systematic Reviews*. 2010;Issue 8. Art. No.: CD006173.
44. Cotter AM, Ness A, Tolosa JE. Prophylactic oxytocin for the third stage of labour. *Cochrane Database of Systematic Reviews*. 2001;Issue 4. Art. No.: CD001808.
45. Peña-Martí G, Comunián-Carrasco G. Fundal pressure versus controlled cord traction as part of the active management of the third stage of labour. *Cochrane Database of Systematic Reviews*. 2007;Issue 4. Art. No.: CD005462.
46. Begley CM, Gyte GML, Murphy DJ, Devane D, McDonald SJ, McGuire W. Active versus expectant management for women in the third stage of labour. *Cochrane Database of Systematic Reviews* 2010;Issue 7. Art. No.: CD007412.
47. McDonald SJ, Abbott JM, Higgins SP. Prophylactic ergometrine-oxytocin versus oxytocin for the third stage of labour. *Cochrane Database of Systematic Reviews*. 2004;Issue 1. Art. No.: CD000201.
48. Gülmezoglu AM, Forna F, Villar J, Hofmeyr GJ. Prostaglandins for preventing postpartum haemorrhage. *Cochrane Database of Systematic Reviews*. 2007;Issue 3. Art. No.: CD000494.
49. McDonald SJ, Middleton P. Effect of timing of umbilical cord clamping of term infants on maternal and neonatal outcomes (Review). *Cochrane Database of Systematic Reviews*. 2008;Issue 2. Art. No.: CD004074.
50. Gülmezoglu AM, Crowther CA, Middleton P. Induction of labour for improving birth outcomes for women at or beyond term. *Cochrane Database for Systematic Reviews*. 2006;Issue 4. Art. No.: CD004945.
51. Mousa HA, Alfirevic Z. Treatment for primary postpartum haemorrhage. *Cochrane Database of Systematic Reviews*. 2007;Issue 1. Art. No.: CD003249.
52. Lopez LM, Hiller JE, Grimes DA. Education for contraceptive use by women after childbirth. *Cochrane Database for Systematic Reviews*. 2010;Issue 1. Art. No.: CD001863.
53. Dodd J, Dare MR, Middleton P. Treatment for women with postpartum iron deficiency anaemia. *Cochrane Database for Systematic Reviews*. 2004;Issue 4. Art. No.: CD004222.
54. French L, Smaill FM. Antibiotic regimens for endometritis after delivery. *Cochrane Database of Systematic Reviews*. 2004;Issue 4. Art. No.: CD001067.
55. McCall EM, Alderdice F, Halliday HL, Jenkins JG, Vohra S. Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants. *Cochrane Database of Systematic Reviews* 2010;Issue 3. Art. No.: CD004210.
56. Imdad A, Yakoob MY, Bhutta ZA. Effect of breastfeeding promotion interventions on breastfeeding rates, with special focus on developing countries. *BMC Public Health* 2011;11(Suppl 3):S24.
57. Dyson L, McCormick FM, Renfrew MJ. Interventions for promoting the initiation of breastfeeding. *Cochrane Database of Systematic Reviews*. 2005;Issue 2. Art. No.: CD001688.
58. Lewin S, Munabi-Babigumira S, Glenton C, et al. Lay health workers in primary and community health care for maternal and child health and the management of infectious diseases. *Cochrane Database of Systematic Reviews*. 2010;Issue 3. Art. No.: CD004015.
59. Lassi ZS, Haider BA, Bhutta ZA. Community-based intervention packages for reducing maternal and neonatal morbidity and mortality and improving neonatal outcomes. *Cochrane Database of Systematic Reviews*. 2010;Issue 11. Art. No.: CD007754.
60. Zupan J, Garner P, Omari AAA. Topical umbilical cord care at birth. *Cochrane Database of Systematic Reviews*. 2004;Issue 3. Art. No.: CD001057.
61. O'Donnell CPF, Davis PG, Morley CJ. Positive end-expiratory pressure for resuscitation of newborn infants at birth. *Cochrane Database of Systematic Reviews*. 2004;Issue 3. Art. No.: CD004341.
62. Grein AJ, Weiner GM. Laryngeal mask airway versus bag-mask ventilation or endotracheal intubation for neonatal resuscitation. *Cochrane Database of Systematic Reviews*. 2005;Issue 2. Art. No.: CD003314.
63. Ziino AJA, Davies MW, Davis PG. Epinephrine for the resuscitation of apparently stillborn or extremely bradycardic newborn infants (Review). *Cochrane Database of Systematic Reviews*. 2002;Issue 3. Art. No.: CD003849.
64. Ungerer RLS, Lincetto O, McGuire W, Saloojee HH, Gülmezoglu AM. Prophylactic versus selective antibiotics for term newborn infants of mothers with risk factors for neonatal infection. *Cochrane Database of Systematic Reviews*. 2004;Issue 4. Art. No.: CD003957.
65. Sazawal S, Black RE. Effect of pneumonia case management on mortality in neonates, infants, and preschool children: a meta-analysis of community based trials. *Lancet Infect Dis*. 2003;3(9):547-556.
66. Anita K M Zaidi, Hammad A Ganatra, Syed S, et al. Effect of case management on neonatal mortality due to sepsis and pneumonia. *BMC Public Health*. 2011;11(Suppl 3):S13.
67. Gordon A, Jeffery HE. Antibiotic regimens for suspected late onset sepsis in newborn infants. *Cochrane Database of Systematic Reviews*. 2005;Issue 3. Art. No.: CD004501.
68. Mtitimila EI, Cooke RWI. Antibiotic regimens for suspected early neonatal sepsis. *Cochrane Database of Systematic Reviews*. 2004;Issue 4. Art. No.: CD004495.
69. Bhutta ZA, Zaidi AK, Thaver D, Humayun Q, Ali S, Darmstadt GL. Management of Newborn Infections in Primary Care Settings: A Review of the Evidence and Implications for Policy? *Pediatric Infectious Disease Journal*. 2009;28(1):S22-30.
70. Lawn JE, Mwansa-Kambafwile J, Horta BL, Barros FC, Cousens S. 'Kangaroo mother care' to prevent neonatal deaths due to preterm birth complications. *International Journal of Epidemiology*. 2010;39(suppl 1):i144 - i154.
71. Conde-Agudelo A, Belizán JM, Diaz-Rossello J. Kangaroo mother care to reduce morbidity and mortality in low birthweight infants. *Cochrane Database of Systematic Reviews*. 2011;Issue 3. Art. No.: CD002771.
72. Edmond K, Bahl R. *Optimal feeding of low-birth-weight infants*. Geneva: World Health Organization; 2006.
73. Soll R. Synthetic surfactant for respiratory distress syndrome in preterm infants. *Cochrane Database of Systematic Reviews*. 1998;Issue 3. Art. No.: CD001149.
74. Greenough A, Dimitriou G, Prendergast M, Milner AD. Synchronized mechanical ventilation for respiratory support in newborn infants. *Cochrane Database of Systematic Reviews*. 2008;Issue 1. Art. No.: CD000456.
75. Ho JJ, Subramaniam P, Henderson-Smart DJ, Davis PG. Continuous distending pressure for respiratory distress syndrome in preterm infants. *Cochrane Database of Systematic Reviews*. 2002;Issue 2. Art. No.: CD002271.
76. Thayyil S, Milligan D. Single versus double volume exchange transfusion in jaundiced newborn infants. *Cochrane Database of Systematic Reviews*. 2006;Issue 4. Art. No.: CD004592.
77. Mills JF, Tudehope D. Fiberoptic phototherapy for neonatal jaundice. *Cochrane Database of Systematic Reviews*. 2001;Issue 1. Art. No.: CD002060.
78. Imdad A, Yakoob MY, Bhutta Z. Effect of breastfeeding promotion interventions on breastfeeding rates, with special focus on developing countries. *BMC Public Health*. 2011;11(Suppl 3):S24.

79. Kramer MS, Kakuma R. Optimal duration of exclusive breastfeeding. *Cochrane Database of Systematic Reviews*. 2002;Issue 1. Art. No.: CD003517.
80. Dewey KG, Adu Afarwuah S. Systematic review of the efficacy and effectiveness of complementary feeding interventions in developing countries. *Maternal & Child Nutrition*. 2008;4(Suppl 1):24-85.
81. Imdad A, Yakoob MY, Bhutta ZA. Impact of maternal education about complementary feeding and provision of complementary foods on child growth in developing countries. *BMC Public Health*. 2011;11(Suppl 3):S25.
82. Eisele TP, Larsen D, Steketee RW. Protective efficacy of interventions for preventing malaria mortality in children in Plasmodium falciparum endemic areas. *International Journal of Epidemiology*. 2010;39(suppl 1):i88.
83. Lengeler C. Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database of Systematic Reviews*. 2004.
84. Thwing J, Eisele TP, Steketee RW. Protective efficacy of malaria case management and intermittent preventive treatment for preventing malaria mortality in children: a systematic review for the Lives Saved Tool. *BMC Public Health*. 2011;11(Suppl 3):S14.
85. Chetty T, Naidu KK, Newell ML. A systematic review of HIV-free survival by feeding practices From birth to 18 months: World Health Organization 2010.
86. Soares-Weiser K, Goldberg E, Tamimi G, Leibovici L, Pitan F. Rotavirus vaccine for preventing diarrhoea. *Cochrane Database of Systematic Reviews*. 2004;Issue 1. Art. No.: CD002848.
87. Soares-Weiser K, Macle hose H, Ben-Aharon I, Goldberg E, Pitan F, Cunliffe N. Vaccines for preventing rotavirus diarrhoea: vaccines in use. *Cochrane Database for Systematic Reviews*. 2010;Issue 5. Art. No.: CD008521.
88. Mayo-Wilson E, Imdad A, Herzer K, Yakoob MY, Bhutta ZA. Vitamin A supplements for preventing mortality, illness, and blindness in children aged under 5: systematic review and meta-analysis. *BMJ*. 2011;343:d5094.
89. Imdad A, Herzer K, Mayo-Wilson E, Yakoob MY, Bhutta ZA. Vitamin A supplementation for preventing morbidity and mortality in children from 6 months to 5 years of age. *Cochrane Database of Systematic Reviews*. 2010;Issue 12. Art. No.: CD008524.
90. Ahmed T, Ahmed SAM, Mahfuz M, Abdullah K, Cravioto A, Sack D. Systematic review of management of childhood severe malnutrition. *Nutrition intervention for maternal and child health and survival*. Vol ISBN- 978-0-19-547360-5. Oxford University Press.; 2010.
91. Community management of severe acute malnutrition: World Health Organization/World Food Programme/United Nations System Standing Committee on Nutrition/The United Nations Children's Fund; 2007.
92. Theodoratou E, Al-Jilaihawi S, Woodward F, et al. The effect of case management on childhood pneumonia mortality in developing countries. *International Journal of Epidemiology*. 2010;39(suppl 1):i155-i171.
93. Fawzi WW, Chalmers TC, Herrera MG, Mosteller F. Vitamin A supplementation and child mortality. A meta-analysis. *JAMA*. 1993;269(7):898-903.
94. Sudfeld CR, Navar AM, Halsey NA. Effectiveness of measles vaccination and vitamin A treatment. *International journal of epidemiology*. 2010;39(suppl 1):i48-i55.
95. Brown N, Roberts C. Vitamin A for acute respiratory infection in developing countries: a meta-analysis. *Acta Paediatrica*. 2004;93(11):1437-1442.
96. Wu T, Ni J, Wei J. Vitamin A for non-measles pneumonia in children. *Cochrane Database of Systematic Reviews*. 2005;Issue 3. Art. No.: CD003700.
97. Grotto I, Mimouni M, Gdalevich M, Mimouni D. Vitamin A supplementation and childhood morbidity from diarrhoea and respiratory infections: A meta-analysis. *Journal of Pediatrics*. 2003;142(3):297-304.
98. Chen H, Zhuo Q, Yuan W, Wang J, Wu T. Vitamin A for preventing acute lower respiratory tract infections in children up to seven years of age. *Cochrane Database of Systematic Reviews*. 2008;Issue 1. Art. No.: CD006090.
99. Yakoob MY, Theodoratou E, Jabeen A, et al. Preventive zinc supplementation in developing countries: impact on mortality and morbidity due to diarrhoea, pneumonia and malaria. *BMC Public Health*. 2011;11(Suppl 3):S23.
100. Lazzarini M, Ronfani L. Oral zinc for treating diarrhoea in children. *Cochrane Database of Systematic Reviews*. 2008;Issue 3. Art. No.: CD005436.
101. Gregorio GV, Gonzales MLM, Dans LF, Martinez EG. Polymer-based oral rehydration solution for treating acute watery diarrhoea. *Cochrane Database of Systematic Reviews*. 2009;Issue 2. Art. No.: CD006519.
102. Hahn S, Kim Y, Garner P. Reduced osmolarity oral rehydration solution for treating dehydration caused by acute diarrhoea in children. *Cochrane Database of Systematic Reviews*. 2002;Issue 1. Art. No.: CD002847.
103. Hartling L, Bellemare S, Wiebe N, Russell KF, Klassen TP, Craig WR. Oral versus intravenous rehydration for treating dehydration due to gastroenteritis in children. *Cochrane Database of Systematic Reviews*. 2006;Issue 3. Art. No.: CD004390.
104. Munos MK, Fischer-Walker CL, Black RE. The effect of rotavirus vaccine on diarrhoea mortality. *International journal of epidemiology*. 2010;39(suppl 1):i56-i62.
105. Walker CLF, Black RE. Zinc for the treatment of diarrhoea: effect on diarrhoea morbidity, mortality and incidence of future episodes. *International Journal of Epidemiology*. 2010;39(suppl 1):i63-i69.
106. Traa BS, Fischer Walker CL, Munos M, Black RE. Antibiotics for the treatment of dysentery in children. *International Journal of Epidemiology*. 2010;39:i70-i74.
107. Christopher PRH, David KV, John SM, Sankarapandian V. Antibiotic therapy for Shigella dysentery. *Cochrane Database of Systematic Reviews*. 2010;Issue 8. Art. No.: CD006784.
108. Gogia S, Sachdev HS. Home visits by community health workers to prevent neonatal deaths in developing countries: a systematic review. *Bulletin of the World Health Organization*. 2010;88(9):658-666.

# ACKNOWLEDGEMENTS



UNDP · UNFPA · WHO · World Bank  
Special Programme of Research, Development  
and Research Training in Human Reproduction



THE AGA KHAN UNIVERSITY









**The Partnership for Maternal, Newborn & Child Health**

World Health Organization

20 Avenue Appia , CH-1211 Geneva 27, Switzerland

Fax: + 41 22 791 5854

Telephone: + 41 22 791 2595

[pmnch@who.int](mailto:pmnch@who.int)

[www.pmnch.org](http://www.pmnch.org)